

Gender disparities in Lyme disease: Women face higher risk of severe and prolonged illness

July 19 2023



Credit: Pixabay/CC0 Public Domain

Women with Lyme disease take longer to get diagnosed, have more severe symptoms and experience higher rates of disability when

compared to men. They may also be more likely to develop persistent Lyme disease. Those are among the findings of a recent study that analyzed information from the MyLymeData patient registry. The results have been published in the *International Journal of General Medicine*.

The present study, which was conducted by LymeDisease.org, a research and advocacy organization, assessed sex-based differences in Lyme disease patients who remained ill for six months or more after antibiotic treatment. In addition to responses from 2,170 patients enrolled in MyLymeData, the researchers reviewed other Lyme disease studies to evaluate the distribution of patients by sex, stage of illness, data source, and enrollment criteria.

Analysis of the study data revealed several important insights. Women reported a higher prevalence of tick-borne co-infections, more [severe symptoms](#), longer diagnostic delays, increased misdiagnoses, and greater functional impairment compared to men. However, no significant differences were observed in antibiotic treatment response or side effects. The majority of both men and women who were treated with antibiotics reported improvement.

Lyme disease is a tick-borne infection caused by *Borrelia burgdorferi*, a corkscrew-shaped bacteria known as a spirochete. The Centers for Disease Control and Prevention (CDC) estimates that over 476,000 people are diagnosed with the illness each year in the United States. That makes new cases of Lyme disease more common than [breast cancer](#), HIV/AIDS and tuberculosis combined in this country.

According to LymeDisease.org CEO Lorraine Johnson, lead author of the study, "We need to find out why more women than men develop persistent Lyme disease. Our study shows that women are diagnosed later—which increases their risk of chronic illness. But we must also determine whether biological variables play a role as they do in other

[infectious diseases."](#)

"Science has ignored gender-based differences in research for far too long," said Dr. Raphael Stricker, the study's senior author. "Our findings underscore the need for greater consideration of gender-specific factors in Lyme disease research in order to improve outcomes for all patients."

More information: Lorraine Johnson et al, Does Biological Sex Matter in Lyme Disease? The Need for Sex-Disaggregated Data in Persistent Illness, *International Journal of General Medicine* (2023). [DOI: 10.2147/IJGM.S406466](https://doi.org/10.2147/IJGM.S406466)

Provided by Taylor & Francis

Citation: Gender disparities in Lyme disease: Women face higher risk of severe and prolonged illness (2023, July 19) retrieved 15 May 2024 from <https://medicalxpress.com/news/2023-07-gender-disparities-lyme-disease-women.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--