

Older adults who identify as sexual and gender minorities experience disproportionate age-related decline, finds study

July 25 2023



Credit: Pixabay/CC0 Public Domain

In June, the U.S. Census Bureau announced that the average age of the U.S. population reached a new high of nearly 39 years old. Today, roughly three million older adults who belong to a sexual and gender minority (individuals who self-describe as LGBTQ+ or as having discordant gender and sex assigned at birth) live in the U.S., a number that is projected to grow to more than five million by 2030.

Prior studies have demonstrated that this demographic is at disproportionate risk of chronic conditions, [mental health issues](#), cognitive decline and health risk behaviors compared to the general population. However, there's little data available to characterize the way this population is aging.

In a new study led by Chelsea N. Wong, MD, an advanced geriatric medicine fellow at Beth Israel Deaconess Medical Center, investigators developed an index to describe and compare frailty—a measure of physiological reserve or vulnerability—between older sexual and gender minority (OSGM) adults and non-OSGM participants.

Using the frailty index they constructed, Wong and colleagues showed that, overall, OSGM had higher rates of frailty, starting at younger ages and continuing into older age, compared to non-OSGM. The team's work appears in the *The Journals of Gerontology: Series A*.

"We found that older individuals who belong to sexual and gender minorities have a higher burden of frailty, maintained across the lifespan and a higher magnitude of mortality risk," said Wong, who is also entering a T32 Post-Doctoral Fellowship with the Harvard Translation Research in Aging Training Program at Hebrew SeniorLife. "To our knowledge this is the first study to examine frailty among this marginalized population."

Frailty is the state of vulnerability caused by age-associated decline across multiple functions, and it is associated with an increased risk of falls, disability, hospitalization, and mortality. While no single symptom confirms a frailty diagnosis, it commonly involves muscle loss and weakness, fatigue, slow walking speed and impaired cognition. Taken together, the symptoms of frailty can lead to [social isolation](#) or make independent living impossible.

Wong and colleagues constructed their frailty index using data from the All of Us Research Program, a program funded by the National Institutes of Health designed to recruit a diverse cohort of participants who have been historically underrepresented in biomedical research. The team built their frailty index to calculate based on 33 factors where individuals may experience deficits, including cognition, cardiovascular diseases, the ability to bathe or run errands alone, maintain a social life, fatigue, anxiety and depression, hearing and sight.

When the researchers applied the Frailty Index to a sample of more than 76,000 eligible All of Us participants, they saw clear patterns. The 9,110 OSGM participants had a higher burden of frailty than their counterparts, with 41% robust (not frail), 34% pre-frail, and 26% frail. In comparison, non-OSGM participants, 50% were robust, 32% were pre-frail and 19% were frail. Further, the risk of mortality associated with frailty was higher among OSGM compared to non-OSGM.

These disparities, the authors note, are understood through the minority stress theory, which describes how the external stressors of discrimination become internalized. Previous research has demonstrated that minority stress is associated with negative health consequences, including changes in inflammation immune and endocrine function, which in turn contribute to poor cardiovascular, metabolic and immunologic clinical outcomes.

"This current generation of [older adults](#) who identify as sexual and gender minorities experienced significant events specific to their minority identity, including homosexuality being officially labeled as a [mental illness](#), the Lavender Scare where there was mass dismissal of SGM workers from U.S. government employment, and the advent of the HIV/AIDS epidemic," said Wong.

"The minority stress experience of chronic additive stress associated

with social stigmatization and discrimination may lead to systematic inflammation and contribute to the development of frailty at relatively younger ages. These findings emphasize the need for early geriatric assessment among OSGM and future work should identify potential targets to mitigate [frailty](#)."

More information: Chelsea N Wong et al, Frailty Among Sexual and Gender Minority Older Adults: The All of Us Database, *The Journals of Gerontology: Series A* (2023). [DOI: 10.1093/gerona/glad149](https://doi.org/10.1093/gerona/glad149)

Provided by Beth Israel Deaconess Medical Center

Citation: Older adults who identify as sexual and gender minorities experience disproportionate age-related decline, finds study (2023, July 25) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-07-older-adults-sexual-gender-minorities.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--