

# Rapid weight loss may improve advanced fatty liver disease, says new research

July 3 2023, by Dimitrios Koutoukidis

---



Credit: AI-generated image ([disclaimer](#))

About 2% of adults worldwide suffer from a condition called [non-alcoholic steatohepatitis](#) (Nash), an advanced form of [non-alcoholic fatty liver disease](#). This occurs when fat builds up in the liver, causing inflammation and scarring.

Without treatment it can eventually lead to [liver cirrhosis](#)—and it can also increase the risk of other serious health conditions, such as heart disease.

There is currently no medication to treat Nash. Since excess fat in the liver is what causes the inflammation and scarring that is characteristic of the condition, the current mainstay treatment for patients is [weight loss](#).

However, the kind of weight loss most people are able to achieve on their own is modest and not enough for significant reductions in liver fat and change inflammation and scarring.

But our recent study, published in the journal *Obesity*, has shown that rapid weight loss achieved through the "[soups and shakes](#)" diet—which is commonly used to treat obesity and type 2 diabetes—may be able to reduce the severity of Nash.

To conduct our study, we recruited 16 participants with obesity, Nash and moderate to advanced liver scarring. Five of the participants were female and 11 were male. Most participants were white.

All of the participants took part in the "soups and shakes" weight loss program, replacing their regular meals with specially formulated soups, shakes and bars for 12 weeks. They consumed four products of their choice daily, which provided them with about 880 calories and all the essential vitamins and minerals.

After the initial 12-week period, they gradually began re-introducing regular food to their diet over the next 12 weeks. They were also given regular support from a dietitian to keep them on track and motivated throughout the 24-week study.

At the start of the study, participants were weighed, had their blood pressure taken, blood tests done and two scans that measured the health of their liver. These scans estimated how advanced their liver inflammation and scarring was and the amount of fat in their liver.

These tests were also repeated at 12 and 24 weeks—with an additional blood test done at four weeks.

Fourteen of the participants completed the 24-week study. Participants lost an average of 15% of their [body weight](#), showing they largely adhered to the weight loss program.

Our study also showed that the rapid weight loss was safe for participants. In the past, this kind of diet program wasn't recommended to Nash patients due to concerns over how safe it may be. The most common side effect patients experienced was constipation—but this was temporary and typically only mild.

Scans also showed that most participants had significant improvements in liver fat and in markers of liver inflammation and scarring.

## **Bigger improvements than medication**

These are some of the largest improvements in liver disease severity [reported in research](#) to date, approaching the level of improvement seen with weight loss after [bariatric surgery](#). [No trialed medication](#) has shown such a large improvement.

While some weight regain is likely to happen, if participants are able to maintain at least most of their [weight](#) loss after the study ends, this could possibly reverse the trajectory of their liver disease.

What's more, systolic [blood pressure](#) and hemoglobin A1C (a marker of

blood sugar control) also significantly improved in participants who'd had hypertension and type 2 diabetes at the start of the study. This may suggest that the program could be used to reduce the risk of [heart disease](#), which is the [most common cause of death](#) in people with Nash.

Because our results are only from a small study, further research is needed to test this program in a larger trial with more diverse participants and a control group. It will also be interesting to see whether this program could be useful for patients suffering with more advanced forms of liver disease—such as [liver cirrhosis](#).

But it is promising to see from our study that the diet appears to be safe for people with Nash and effective in improving their liver health.

**More information:** Dimitrios A. Koutoukidis et al, A low-energy total diet replacement program demonstrates a favorable safety profile and improves liver disease severity in nonalcoholic steatohepatitis, *Obesity* (2023). [DOI: 10.1002/oby.23793](https://doi.org/10.1002/oby.23793)

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Rapid weight loss may improve advanced fatty liver disease, says new research (2023, July 3) retrieved 21 May 2024 from <https://medicalxpress.com/news/2023-07-rapid-weight-loss-advanced-fatty.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---