

## New smaller, portable MRI systems could expand delivery of health care, capabilities of medicine

July 17 2023



NIST researchers Kalina Jordanova and Stephen Ogier conduct MRI measurements using a magnetic field strength of 64 millitesla (mT), which is at least 20 times lower than the magnetic field in conventional MRI scanners, used for example in hospital settings. Credit: R. Jacobson/NIST

Magnetic resonance imaging (MRI) machines can clearly view non-bony



parts of the body—soft tissue such as the brain, muscles and ligaments—as well as detect tumors, making it possible to diagnose many diseases and other conditions. However, the powerful magnets in conventional MRI machines make them expensive and bulky, confining them mainly to hospitals and other large facilities.

As an alternative solution, companies are developing new portable versions that have lower-strength magnetic fields. These new models can potentially expand the ways in which MRI is used. For instance, lowfield MRI systems could be deployed in ambulances and other mobile settings. They also could cost much less, promising to make MRI more widely available, including in underserved communities and developing nations.

But for low-field MRI scanners to reach their full potential, more research is needed to understand the relationship between low-field images and the underlying tissue properties they represent. Researchers at the National Institute of Standards and Technology (NIST) have been working on several fronts to advance low-field MRI technology and validate methods for creating images with weaker magnetic fields.

"Magnetic resonance images of tissue differ depending on magnetic strength," said NIST electrical engineer Kalina Jordanova. "With lowfield MRI systems, the contrast of the images is different, so we need to know how human tissue looks at these lower field strengths."





Kalina Jordanova, Stephen Ogier and Katy Keenan are among the NIST researchers who have been working on several projects that aim to advance MRI technology that uses lower-strength magnetic fields and validate its approaches for capturing images with weaker magnetic fields. Credit: R. Jacobson/NIST

Toward these ends, researchers measured the properties of brain tissue at low <u>magnetic field</u> strength. Their results were published in the journal *Magnetic Resonance Materials in Physics, Biology and Medicine.* 

The researchers used a commercially available portable MRI machine to image brain tissue in five male and five female volunteers. The images were created using a magnetic field strength of 64 millitesla, which is at least 20 times lower than the magnetic field in conventional MRI scanners.

The team collected images of the entire brain and obtained data on its



gray matter (which has a high concentration of nerve cells), white matter (deeper tissues of the brain that house nerve fibers), and cerebrospinal fluid (clear fluid surrounding the brain and spinal cord).

These three <u>brain</u> constituents respond to the low magnetic field in different ways and produce distinctive signals that reflect their unique properties, enabling the MRI system to produce images that contain quantitative information about each constituent. "Knowing the quantitative properties of tissue allows us to develop new image collection strategies for this MRI system," said NIST biomedical engineer Katy Keenan.



NIST researcher Sam Oberdick explored contrasts agents for MRI machines along with his cohort. His group tested iron oxide nanoparticles at lower-strength magnetic fields. These nanoparticles inside the liquid solution (pictured here) are



magnetic and are pulled toward the magnet through a combination of magnetic interactions and surface tension. Credit: R. Wilson/NIST

In separate work published in *Scientific Reports*, NIST researchers collaborated with the University of Florence in Italy and Hyperfine Inc. in Guilford, Connecticut, to explore several candidate materials that can significantly boost image quality in low-field MRI scans.

MRI contrast agents—magnetic materials that are injected into patients and enhance image contrast—make it easier for radiologists to identify anatomical features or evidence of disease and are routinely used in MRI at conventional magnetic field strengths. However, researchers are just starting to understand how contrast agents might be used with the new low-field MRI scanners. At the lower field strengths of these scanners, contrast agents may act differently than at higher field strengths, creating opportunities to use new types of magnetic materials for image enhancement.

NIST scientists and their colleagues compared the sensitivity of several magnetic contrast agents in low magnetic fields. The researchers found that <u>iron oxide nanoparticles</u> outperformed traditional contrast agents, which are made of the element gadolinium—a rare-earth metal. At low magnetic field strength, the nanoparticles provided good contrast using a concentration of only about one-ninth that of the gadolinium particles.

Iron oxide nanoparticles also offer the advantage that they are broken down by the human body instead of potentially accumulating in tissue, noted NIST researcher Samuel Oberdick. By comparison, a small amount of gadolinium may accumulate in tissue and could confound the interpretation of future MRI scans if it is not taken into account.



**More information:** Kalina V. Jordanova et al, In vivo quantitative MRI: T1 and T2 measurements of the human brain at 0.064 T, *Magnetic Resonance Materials in Physics, Biology and Medicine* (2023). DOI: 10.1007/s10334-023-01095-x

Samuel D. Oberdick et al, Iron oxide nanoparticles as positive T1 contrast agents for low-field magnetic resonance imaging at 64 mT, *Scientific Reports* (2023). DOI: 10.1038/s41598-023-38222-6

Provided by National Institute of Standards and Technology

Citation: New smaller, portable MRI systems could expand delivery of health care, capabilities of medicine (2023, July 17) retrieved 9 May 2024 from https://medicalxpress.com/news/2023-07-smaller-portable-mri-delivery-health.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.