

Study finds younger kidney cancer survivors at significant risk for heart problems

July 5 2023



Credit: Pixabay/CC0 Public Domain

New research out of VCU Massey Cancer Center indicates that many adolescent and young adult kidney cancer survivors are at a significantly elevated risk for heart issues.

Cardiovascular disease is a leading cause of health complications and

death among adolescents and young adults (AYAs) diagnosed with cancer, where AYAs are characterized as patients between the ages of 15 and 39.

A study published July 5 in the *Journal of the National Comprehensive Cancer Network* assessed the incidence and risk of hypertension—high blood pressure—and [heart failure](#) among AYA patients diagnosed with kidney cancer who also received a form of drug that blocks [blood vessel growth](#) as part of their treatment.

In particular, the researchers looked at the effects of two drugs called sunitinib and sorafenib. They found that approximately half of AYAs treated with sorafenib and one-third of AYAs treated with sunitinib eventually developed hypertension.

"The large number of AYAs who had [high blood pressure](#) during treatment with sunitinib or sorafenib suggests that even individuals without identifiable pre-existing factors—such as older age, obesity and male gender—are also at significant risk for hypertension from these drugs," said study lead author Wendy Bottinor, M.D, a cardio-oncologist and member of the Cancer Prevention and Control research program at Massey and VCU Health Pauley Heart Center.

Contrary to their original hypothesis, the researchers discovered that younger age was not associated with a reduced risk of heart failure in AYA cancer survivors compared to older cancer patients. In fact, this population is at risk for a type of heart failure called left ventricular systolic dysfunction.

In the United States, approximately 90,000 AYAs are diagnosed with cancer every year, [according to the American Cancer Society](#). Kidney, thyroid and colorectal tumors are among the more common cancers in this age group, a trend that has been on the rise over the last few

decades.

The risk for [heart disease](#) among AYAs with cancer is more than double that of people in the same age group without cancer, and the risk of death is nearly 10 times higher among AYAs with heart disease when compared with AYAs who do not have heart disease, as indicated by multiple studies over the last decade. Hypertension forces the heart and blood vessels to work overtime, eventually damaging the tissues in the arteries and increasing a person's odds of irregular heartbeat, heart attack or stroke.

Chemical signals within the body regulate a process called angiogenesis, which is the production of new blood vessels. One of these chemical signals—[vascular endothelial growth factor](#) (VEGF)—clings to the surface of other cells to influence the growth and survival of new blood vessels.

Angiogenesis plays a critical role in the progression of solid tumors because cancer cells need the oxygen and nutrients from blood to grow and spread. A class of drugs called angiogenesis inhibitors are often used alone or in combination with other therapies to stymie the growth of blood vessels that support tumor growth.

This study looked at early-stage [kidney cancer](#) patients who received a specific type of angiogenesis inhibitor—VEGF inhibitors—as part of their treatment regimen. Sunitinib and sorafenib are VEGF inhibitors.

"Although VEGF inhibitors are often used as an effective therapeutic option for adult and pediatric cancer patients, cardiovascular toxicities can be a significant limitation of this treatment, with hypertension and left ventricular dysfunction among the most common," Bottinor said. Historically, the scientific understanding of the cardiovascular toxicities of these drugs among AYAs has been very limited.

"Adolescent and [young adults](#) are an underrepresented group in cancer research with a significant cardiovascular burden," Bottinor said.

"Understanding the relationship between cancer diagnosis, treatment and [heart](#) disease is imperative to promoting cardiovascular health over the entire lifetime of adolescent and young adult [cancer](#) survivors."

Findings from this study warrant a need for further research to understand and reduce the factors that influence cardiovascular risk in this population.

Provided by Virginia Commonwealth University

Citation: Study finds younger kidney cancer survivors at significant risk for heart problems (2023, July 5) retrieved 9 May 2024 from <https://medicalxpress.com/news/2023-07-younger-kidney-cancer-survivors-significant.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--