

# Outdoor air pollution may increase non-lung cancer risk in older adults

August 3 2023

---



Credit: Unsplash/CC0 Public Domain

Chronic exposure to fine particulate air pollutants (PM<sub>2.5</sub>) and nitrogen dioxide (NO<sub>2</sub>) may increase non-lung cancer risk in older adults, according to a study led by Harvard T.H. Chan School of Public Health.

In a cohort study of millions of Medicare beneficiaries, the researchers found that exposures to PM<sub>2.5</sub> and NO<sub>2</sub> over a 10-year period increased the risk of developing colorectal and prostate cancers. The researchers also found that even low levels of air pollution exposure may make people particularly susceptible to developing these cancers, in addition to breast and endometrial cancers.

"Our findings uncover the biological plausibility of air pollution as a crucial risk factor in the development of specific cancers, bringing us one step closer to understanding the impact of air pollution on human health," said Yaguang Wei, research fellow in the Department of Environmental Health. "To ensure equitable access to clean air for all populations, we must fully define the effects of air pollution and then work towards reducing it."

The study was published online August 1, 2023, in *Environmental Epidemiology*.

While air pollution has been established as a risk factor for lung cancer, and a link to breast cancer risk has been emerging, few studies have looked at its effects on prostate, colorectal, and endometrial cancer risk.

Researchers analyzed data from national Medicare beneficiaries aged 65 or older, collected from 2000 to 2016. All subjects were cancer-free for at least the initial 10 years of the study period. The researchers created separate cohorts for each type of cancer—breast, colorectal, endometrial, and prostate—with between 2.2 million and 6.5 million subjects in each cohort. Separate analyses looked at cancer risk under the impacts of air pollutants for various subgroups by factors including age, sex (for colorectal cancer only), race/ethnicity, average BMI, and socioeconomic status.

Drawing from a variety of air pollution data sources, the researchers

developed a predictive map of PM<sub>2.5</sub> and NO<sub>2</sub> concentrations across the contiguous U.S. This was then linked to beneficiaries' residential ZIP codes to enable the researchers to estimate individual exposures over a 10-year period.

Findings from the nationwide analysis showed that chronic PM<sub>2.5</sub> and NO<sub>2</sub> exposures increased the risk of developing colorectal and prostate cancers but were not associated with endometrial cancer risk. For breast cancer, NO<sub>2</sub> exposure was associated with an increased risk, while the association for PM<sub>2.5</sub> was inconclusive. The researchers suggested that the mixed associations may be due to variations in the chemical composition of PM<sub>2.5</sub>, which is a complex mixture of solid and liquid particles.

When the analysis was restricted to regions where air pollution levels were significantly below national standards and the composition of PM<sub>2.5</sub> remained fairly stable, their effect on breast cancer risk was more pronounced. Stronger associations between exposures to both pollutants and endometrial cancer risk were also found at lower pollution levels.

[In their analysis of risk by subgroups, the researchers found evidence suggesting that communities with higher average BMI may face disproportionately higher risk of all four cancers from NO<sub>2</sub> exposure, and that Black Americans and those enrolled in Medicaid may be more susceptible to cancer risks \(prostate and breast, respectively\) from PM<sub>2.5</sub> exposure.](#)

The researchers noted that even communities with seemingly clean air were not immune to cancer risk. They found substantial associations between exposure to the two pollutants and the risks of all four cancers even at pollution levels below newly updated World Health Organization guidelines (which are lower than current U.S. standards).

"The key message here is that U.S. air pollution standards are inadequate in protecting public health," said senior author Joel Schwartz, professor of environmental epidemiology. "The Environmental Protection Agency recently proposed stricter standards for PM<sub>2.5</sub>, but their proposal doesn't go far enough in regulating this pollutant. Current NO<sub>2</sub> standards are also woefully inadequate. Unless all of these standards become much, much stricter, air pollution will continue to result in thousands of unnecessary cases of multiple cancers each year."

Other Harvard Chan School authors include Edgar Castro, Cristina Su Liu, Xinye Qiu, James Healy, and Bryan Vu.

---

Correction note: Due to a mistake in interpretation of data, the previous version of the article stated the study found that NO<sub>2</sub> exposure is associated with a decreased risk of breast cancer. The authors have confirmed that the results showed that NO<sub>2</sub> exposure is associated with an increased risk of breast cancer.

**More information:** Yaguang Wei et al, Additive effects of 10-year exposures to PM<sub>2.5</sub> and NO<sub>2</sub> and primary cancer incidence in American older adults, *Environmental Epidemiology* (2023). [DOI: 10.1097/EE9.0000000000000265](https://doi.org/10.1097/EE9.0000000000000265)

Provided by Harvard T.H. Chan School of Public Health

Citation: Outdoor air pollution may increase non-lung cancer risk in older adults (2023, August 3) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-08-outdoor-air-pollution-non-lung-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.