

# Scientists develop finger sweat test to detect antipsychotic drugs in patients

August 31 2023

---



Credit: Pixabay/CC0 Public Domain

Antipsychotic drugs treat incredibly vulnerable patients. Maintaining a treatment regimen is difficult for many patients, but not taking the medication is associated with a higher risk of poor health outcomes. These drugs are also very powerful with strong side effects, and blood tests are often used to calibrate a patient's dosage and confirm that they are taking the recommended dose.

However, blood tests are invasive and potentially uncomfortable. Scientists have now discovered a way to test the levels of common [antipsychotic drugs](#) in the sweat from patients' fingerprints, offering a quicker, more comfortable, and more convenient alternative to blood draws for patient monitoring.

"Our test offers patients a quick and dignified way of showing commitment to antipsychotic treatment," said Katherine Longman of the University of Surrey, first author of the study in *Frontiers in Chemistry*. "This non-invasive approach can also be adapted to fit other therapeutic regimes."

## A test at your fingertips

The scientists already knew that some drugs could be detected in the sweat from a fingertip, without the requirement for specialist personnel and with easier storage and transportation. Finger sweat samples, unlike blood, can be transported at ambient temperature.

To test whether antipsychotic drugs could also be detected in sweat, they recruited 60 patients receiving clozapine, quetiapine, or olanzapine, as well as 30 negative controls. Eleven patients taking clozapine also agreed to supply [blood samples](#) so that the correlation between finger sweat indicators and blood indicators could be tested. Patients were asked to report their dosage and the most recent dose they had taken.

The researchers collected samples both before and after handwashing, as hands that have been washed are considered to give a better picture of the eccrine sweat that comes from fingertips. Patients pressed their fingertips against porous paper for 30 seconds. These samples were then collected and analyzed using liquid chromatography mass spectrometry.

Corresponding author Prof. Melanie Bailey, based at the University of

Surrey, and the team also asked six people who did not take the drugs to handle whole and crushed pills and then give fingerprints. This control checked whether the test could be compromised by patients touching the medication. The scientists were able to confirm the reliability of the method and to distinguish between the presence of the drugs in patients' sweat and the presence of the [drug](#) on patients' hands.

## Fingerprinting medication

The test detected the presence of antipsychotic drugs accurately in every patient taking them. It was most effective for clozapine, where a pilot analysis of a subgroup of patients found that the levels of clozapine metabolites in finger sweat correlated with the levels found in blood. This, the researchers said, raises the exciting possibility that the test will eventually be able to quantify the levels of clozapine in a patient's [sweat](#) instead of just detecting them.

The test also consistently detected quetiapine, although this was a small group of patients. The signal for olanzapine was not as strong, but all patients who took olanzapine tested positive on at least one fingerprint provided before washing their hands. They were also taking lower doses than patients on the other drugs.

Although the test as used in the study included samples taken after washing hands, which adds time and facilities needed for the test, the efficacy of the tests on fingerprints taken from unwashed hands suggested this wasn't necessary. Removing this step would make it even quicker and easier for laypeople to carry out the tests.

"We are currently exploring methods to quantify the level of drug in a fingerprint and the optimum sampling time," added Bailey. "We are also very interested to see whether fingerprints can be used to diagnose disease—for example, from the metabolites that are deposited in a

fingerprint sample."

**More information:** Noninvasive drug adherence monitoring of antipsychotic patients via finger sweat testing, *Frontiers in Chemistry* (2023). [DOI: 10.3389/fchem.2023.1245089](https://doi.org/10.3389/fchem.2023.1245089)

Provided by Frontiers

Citation: Scientists develop finger sweat test to detect antipsychotic drugs in patients (2023, August 31) retrieved 9 May 2024 from <https://medicalxpress.com/news/2023-08-scientists-finger-antipsychotic-drugs-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--