

## Subcutaneous securement system best for reaching end of PICC need

August 31 2023, by Elana Gotkine



A subcutaneous anchor securement system (SASS) is superior for



assisting oncology patients to reach the end of need with a single peripherally inserted central catheter (PICC), according to a study published online Aug. 1 in *The Journal of Vascular Access*.

Michelle L. Hawes, R.N., from Data to Wisdom Research Consulting in Bloomington, Indiana, and colleagues conducted a <u>retrospective study</u> to examine <u>oncology</u> patients' ability to finish their therapy with one PICC using implant and explant data for adult <u>oncology patients</u> from 2007 to 2021. All patients received a PICC with an adhesive securement device (ASD) or SASS.

The researchers found that partial or complete dislodgement causing unplanned PICC removal occurred for 12 and 0.4% of those with ASD and SASS, respectively. The probability of reaching the end of need with one PICC at two years was 68% and more than 95%, respectively, regardless of the reason for premature removal, with a significant difference between the two securement devices (P

"Based on the evidence, the best instrument for securing a catheter throughout a patient's therapeutic need is the subcutaneous anchoring securement device," Hawes said in a statement.

**More information:** Michelle L Hawes et al, A retrospective study of subcutaneous anchor securement systems in oncology patients, *The Journal of Vascular Access* (2023). DOI: 10.1177/11297298231190416

2023 HealthDay. All rights reserved.

Citation: Subcutaneous securement system best for reaching end of PICC need (2023, August 31) retrieved 15 May 2024 from https://medicalxpress.com/news/2023-08-subcutaneous-picc.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.