

Study finds women who consumed sugar sweetened beverage daily had higher risk of developing liver issues

August 8 2023



Credit: CC0 Public Domain

Approximately 65% of adults in the United States consume sugar sweetened beverages daily. Chronic liver disease is a major cause of morbidity and mortality worldwide and can result in liver cancer and

liver disease-related mortality.

Researchers from Brigham and Women's Hospital, a founding member of the Mass General Brigham health care system, led one of the first studies to look at the association between intake of [sugar-sweetened beverages](#), artificially sweetened beverages, and incidence of [liver cancer](#) and [chronic liver disease](#) mortality. Results are published in *JAMA*.

"To our knowledge, this is the first study to report an association between sugar sweetened beverage intake and chronic liver disease mortality," said first author Longgang Zhao, Ph.D., of the Brigham's Channing Division of Network Medicine. Zhao is a postdoctoral researcher who works with senior author Xuehong Zhang, MBBS, ScD, in the Channing Division. "Our findings, if confirmed, may pave the way to a public health strategy to reduce risk of liver disease based on data from a large and geographically diverse cohort."

This [observational study](#) included nearly 100,000 [postmenopausal women](#) from the large, prospective Women's Health Initiative study. Participants reported their usual soft drink, fruit drink (not including fruit juice) consumption, and then reported artificially sweetened beverage consumption after three years.

Participants were followed for a median of more than 20 years. Researchers looked at self-reported liver cancer incidence and death due to chronic liver disease such as fibrosis, cirrhosis, or chronic hepatitis, which were further verified by medical records or the National Death Index.

A total of 98,786 postmenopausal women were included in the final analyses. The 6.8 percent of women who consumed one or more sugar-sweetened beverages daily had an 85 percent higher risk of liver cancer

and 68 percent higher risk of chronic liver disease mortality compared to those who had fewer than three sugar sweetened beverages per month.

The authors note that the study was observational, and causality cannot be inferred, and relied on self-reported responses about intake, sugar content and outcomes. More studies are needed to validate this risk association and determine why the sugary drinks appeared to increase risk of [liver cancer](#) and disease.

Furthermore, more research is needed to elucidate the potential mechanisms by integrating genetics, preclinical and experimental studies, and -omics data.

More information: Longgang Zhao et al, Sugar-Sweetened and Artificially Sweetened Beverages and Risk of Liver Cancer and Chronic Liver Disease Mortality, *JAMA* (2023). [DOI: 10.1001/jama.2023.12618](https://doi.org/10.1001/jama.2023.12618)

Provided by Mass General Brigham

Citation: Study finds women who consumed sugar sweetened beverage daily had higher risk of developing liver issues (2023, August 8) retrieved 21 May 2024 from <https://medicalxpress.com/news/2023-08-women-consumed-sugar-sweetened-beverage.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--