


CT test simulates blood flow to assess risk in patients with angina

September 12 2023



Coronary CT angiography (CTA) images (left side, A and B) and corresponding derived fractional flow reserve (FFR) (right side, A and B). The location 2-cm distal to stenosis is manually delineated and the coronary CTA-derived FFR (FFR_{CT}) value is registered. The lowest lesion-specific coronary CTA-derived FFR value is used to categorize participants, with (A) greater than 0.80 representing a normal value and (B) 0.80 or less representing an abnormal value. Credit: Radiological Society of North America

An advanced CT test can identify individuals with stable angina at a reduced risk of three-year adverse outcomes despite their having a high coronary artery calcium score, according to a study published in the journal *Radiology*.

In the prospective study, researchers in Denmark assessed the three-year clinical outcomes of patients who underwent CT angiography and CT angiography-derived fractional flow reserve (CT-FFR). CT-FFR uses coronary CT angiography images of the heart, AI algorithms and [computational fluid dynamics](#) to simulate the amount of blood flowing through the coronary arteries.

"CT angiography is the first step in identifying the presence of coronary artery disease, but it doesn't show the impact on blood flow in the arteries," said lead researcher Kristian T. Madsen, M.D., and Ph.D. candidate in the Department of Cardiology at the University Hospital of Southern Denmark in Esbjerg. "Turbulent blood flow may create abnormal pressure in the vessel that grinds down coronary plaques making them prone to rupture."

Non-contrast enhanced cardiac CT is typically performed prior to CT angiography to determine the amount of coronary artery [calcium](#), which is a surrogate marker for total coronary plaque, a fatty buildup in the arteries, that can lead to a future heart attack. A coronary artery calcium score of 400 or higher indicates high plaque levels. As calcium absorbs a lot of X-rays, it can be especially challenging to assess coronary CT images of patients with high amounts of coronary artery calcium.

The [study group](#) included patients with new-onset stable coronary artery disease consecutively enrolled in the Assessing Diagnostic Value of Non-Invasive CT-FFR in Coronary Care (ADVANCE) clinical trial between

December 2015 and October 2017 at three Danish sites. Eligibility factors for the trial included at least one coronary artery stenosis, or narrowing, greater than 30% and no [atrial fibrillation](#) ([irregular heartbeat](#)) or prior coronary revascularization, which is a procedure or surgery that improves blood flow to the heart.

Of the 900 participants, 523 had normal CT-FFR results (mean age 64, 318 males) and 377 had abnormal CT-FFR results (mean age 65, 264 males). In the normal results group, the rate of three-year adverse outcomes (defined as all-cause deaths and non-fatal myocardial infarction) was 2.1% (11/523). In the abnormal results group, the rate of three-year adverse outcomes was 6.6% (25/377).

Among the participants with normal CT-FFR results and a high coronary artery calcium score, the three-year adverse outcome rate was 2.2% (4/182). Participants with a high coronary artery calcium score and abnormal CT-FFR results had a 9% (19/212) three-year adverse outcome rate.

"Our study provides evidence for the prognostic potential of CT-FFR in patients with high coronary artery calcium scores," he said. "Regardless of the patient's baseline risk and extent of coronary artery disease measured by coronary artery calcium, if CT-FFR results are normal, the prognosis is good."

According to Dr. Madsen, imaging options to assess the impact of coronary artery disease on blood-flow have different strengths and weaknesses. He said CT-FFR exhibits high diagnostic performance, and its results correlate well with invasively measured fractional flow reserve, an invasive procedure that measures blood flow directly within the coronary arteries.

"In my opinion, CT-FFR is the best option because it provides a [blood-](#)

[flow](#) estimate similar to what you would obtain if you performed invasive heart catheterization," he said. "You get a reliable and quick answer without performing additional tests or putting the patient through an invasive procedure."

Dr. Madsen said CT-FFR will allow clinicians to divide the coronary artery disease patient population by risk level and focus on improving the prognosis of high-risk patients.

"CT-FFR provides diagnostic and prognostic information beyond what can be obtained from CT angiography alone across a wide range of coronary artery calcium levels," he said. "It's a tool that offers a lot of exciting prospects for the future."

More information: Kristian T. Madsen et al, Prognostic Value of Coronary CT Angiography–derived Fractional Flow Reserve on 3-year Outcomes in Patients with Stable Angina, *Radiology* (2023). DOI: 10.1148/radiol.230524 , [dx.doi.org/10.1148/radiol.230524](https://doi.org/10.1148/radiol.230524)

Valentin Sinitsyn, Insights into CT-derived Coronary Fractional Flow Reserve and Coronary Artery Calcification, *Radiology* (2023). DOI: 10.1148/radiol.232150 , [dx.doi.org/10.1148/radiol.232150](https://doi.org/10.1148/radiol.232150)

Provided by Radiological Society of North America

Citation: CT test simulates blood flow to assess risk in patients with angina (2023, September 12) retrieved 27 April 2024 from <https://medicalxpress.com/news/2023-09-ct-simulates-blood-patients-angina.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.