

## **Increasing steps by 3,000 per day can lower blood pressure in older adults**

September 26 2023, by Anna Zarra Aldrich



Change in blood pressure in those taking and not taking anti-hypertensive medications. In participants taking anti-hypertensive medications (n = 8) at baseline, 10 weeks, and 20 weeks, systolic and diastolic blood pressure decreased from  $135 \pm 8$  to  $130 \pm 12$  to  $126 \pm 10$  mm Hg and  $80 \pm 8$  to  $77 \pm 9$  to  $77 \pm 5$  mm Hg, respectively. In participants not on anti-hypertensive medications (n = 13) at baseline, 10 weeks, and 20 weeks, systolic and diastolic blood pressure decreased from  $137 \pm 11$  to  $134 \pm 14$  to  $132 \pm 12$  mm Hg and  $82 \pm 5$  to  $80 \pm 7$  to  $78 \pm 7$  mm Hg, respectively. Error bars indicate standard deviations. Credit: *Journal of Cardiovascular Development and Disease* (2023). DOI: 10.3390/jcdd10080317



An estimated 80% of older adults in the U.S. have high blood pressure. Maintaining healthy blood pressure can protect against serious conditions like heart failure, heart attacks, and strokes.

A new study including Linda Pescatello, distinguished professor of kinesiology in the College of Agriculture, Health and Natural Resources, found that adding a relatively minimal amount of movement, about 3,000 steps per day, can significantly reduce high <u>blood pressure</u> in older adults.

Pescatello worked with Elizabeth Lefferts, the lead author of the paper, Duck-chun Lee, and others in Lee's lab at Iowa State University. They published their findings in a recent issue of the *Journal of Cardiovascular Development and Disease*.

"We'll all get high blood pressure if we live long enough, at least in this country," Pescatello says. "That's how prevalent it is."

Pescatello is an expert on hypertension (the clinical term for <u>high blood</u> <u>pressure</u>) and exercise. Her previous research has demonstrated that exercise can have a significant immediate and long-lasting impact on lowering blood pressure in hypertensive adults.

This study sought to determine if older adults with hypertension could receive these benefits by moderately increasing their daily walking, which is one of the easiest and most popular forms of physical activity for this population.

"It's easy to do, they don't need any equipment, they can do it anywhere at almost any time," Lee says.

The study focused on a group of sedentary <u>older adults</u> between ages 68 and 78 who walked an average of about 4,000 steps per day before the



study.

After consulting existing studies, Lee determined that 3,000 steps would be a reasonable goal. This would also put most participants at 7,000 daily steps, in line with the American College of Sports Medicine's recommendation.

"3,000 steps is large enough but not too challenging to achieve for <u>health</u> <u>benefits</u>," Lee says.

The team conducted the study during the height of the COVID-19 pandemic, which meant they had to do everything remotely.

The researchers sent participants a kit with pedometers, blood pressure monitors, and step diaries for participants to log how much they were walking each day.

On average, participants' systolic and <u>diastolic blood pressure</u> decreased by an average of seven and four points, respectively, after the intervention.

Other studies suggest decreases of these magnitudes correspond to a relative risk reduction of all-cause mortality by 11%, and 16% for cardiovascular mortality, an 18% reduction in the risk of heart disease, and a 36% risk reduction of stroke.

"It's exciting that a simple lifestyle intervention can be just as effective as structured exercise and some medications," Lefferts says.

The findings suggest that the 7,000-step regimen the participants in the study achieved is on-par with reductions seen with anti-hypertensive medications. Eight of the 21 participants were already on anti-hypertensive medications. Those participants still saw improvements in



systolic blood pressure from increasing their daily activity.

"In a <u>previous study</u>, we found that when exercise is combined with <u>medication</u>, exercise bolsters the effects of blood pressure medication alone," Pescatello says. "It just speaks to the value of <u>exercise</u> as anti-hypertensive therapy. It's not to negate the effects of medication at all, but it's part of the treatment arsenal."

The researchers found that walking speed and walking in continuous bouts did not matter as much as simply increasing total steps.

"We saw that the volume of physical activity is what's really important here, not the intensity," Pescatello says. "Using the volume as a target, whatever fits in and whatever works conveys health benefits."

**More information:** Elizabeth C. Lefferts et al, Increasing Lifestyle Walking by 3000 Steps per Day Reduces Blood Pressure in Sedentary Older Adults with Hypertension: Results from an e-Health Pilot Study, *Journal of Cardiovascular Development and Disease* (2023). DOI: 10.3390/jcdd10080317

Provided by University of Connecticut

Citation: Increasing steps by 3,000 per day can lower blood pressure in older adults (2023, September 26) retrieved 11 May 2024 from <u>https://medicalxpress.com/news/2023-09-day-blood-pressure-older-adults.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.