


Death rates after first heart attack have gone down for those without diabetes or with type 2 diabetes, but not type 1

September 12 2023



Credit: CC0 Public Domain

New research to be presented at this year's [Annual Meeting of the European Association for the Study of Diabetes \(EASD\)](#) in Hamburg,

Germany (2–6 October) shows that following a heart attack, there have been falls in the death rates of both people without diabetes and those with type 2 diabetes, but not those with type 1 diabetes. The study is by Dr. Linn Glynn, Karolinska Institutet, Stockholm, Sweden, and colleagues.

Earlier studies have demonstrated that people with diabetes have a worse outcome after [heart](#) attacks ([myocardial infarction](#) [MI]), compared to people without diabetes. In contrast, recent studies have shown a substantially decline in overall mortality and cardiovascular (CV) outcome among people with diabetes. However, trends in survival after a first [heart attack](#) in people with diabetes and without diabetes have still not been extensively studied.

Using data from between 2006 to 2020, the authors identified 2,527 individuals with type 1 diabetes (T1D), 48,321 individuals with type 2 diabetes (T2D) and 243,170 individuals without diabetes with a [first heart attack](#) in national health care registries. The outcomes measured were trends in survival, (overall mortality and cardiovascular [CV] [death](#)), and major cardiovascular events (MACE), meaning the combined outcomes of non-fatal stroke, non-fatal MI, CV death and hospitalized heart failure. Computer and statistical modeling was used to establish any variations in these outcomes.

Individuals with T1D were younger (62 years) and more often women (44%) compared to individuals with T2D (75 years,) women (38%), and to the [control group](#) without diabetes (73 years) women (38%). Three-year trends in death, CV death and MACE between groups are shown in the full abstract.

During follow-up and after multiple adjustments (sex, age, comorbidities, socioeconomic factors and medication) there was a significant decreased annual incidence trend for all cause death in

control group (-1.9% per year) and individuals with T2D (-1.3% per year) with no such trend in individuals with T1D (no change). Corresponding numbers for CV death were for control group -2.0% per year, T2D -1.6% per year, and T1D -0.5% per year, but the result for T1D was not statistically significant. For MACE, incidence in the control group fell by 2.3% per year, T2D by 1.9% per year and T1D by 0.6% per year, but again the result for T1D was not statistically significant.

Discussing potential reasons for the findings, the authors say that the standard care following a heart attack has improved with more availability of—for example—percutaneous coronary intervention and better overall medical treatment. However, this standard of care should have improved in all three groups.

In this study population, the authors say that people with T1D had a longer duration of diabetes and a higher mean glycated hemoglobin (a measure of the average blood sugar level over the past 3 months) than people with T2D, which might have an impact on the risk of developing a heart attack as well as the prognosis following a heart attack.

The authors conclude, "During the last 15 years, the risk of death and major cardiovascular events in people without diabetes and with type 2 diabetes after having a first-time heart attack has decreased significantly. In contrast, this decreasing trend was absent in people with type 1 [diabetes](#). Our study highlights the urgent need for understanding the cardiovascular risk in people with T1D."

Provided by Diabetologia

Citation: Death rates after first heart attack have gone down for those without diabetes or with type 2 diabetes, but not type 1 (2023, September 12) retrieved 2 May 2024 from <https://medicalxpress.com/news/2023-09-death-heart-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.