

Exploring durability of hepatitis B surface antigen seroclearance using real-world data from electronic health records

September 19 2023



Patient inclusion and exclusion flowchart. HBsAg, hepatitis B surface; HBV, hepatitis B virus; IFN, interferon; NAs, nucleos(t)ide analogue. Credit: *Genes & Diseases*



In a <u>study</u> published in the journal *Genes & Diseases*, researchers from The Second Affiliated Hospital of Chongqing Medical University analyzed data from an extensive dataset comprising over 70,000 HBsAgpositive individuals at The Second Affiliated Hospital of Chongqing Medical University.

They compared two groups: those achieving HBsAg seroclearance through NAs <u>monotherapy</u> (168 patients) and those through IFN monotherapy (30 patients). NAs monotherapy patients were older, with a higher proportion achieving HBsAg seroclearance during treatment (48.8% vs. 30.0%). They also had a more favorable immune response, with a higher percentage developing concomitant anti-HBs.

The NAs-monotherapy group had more individuals with normal ALT levels (78.6% vs. 40.0%). The time to achieve HBsAg seroclearance and therapy duration were similar between the groups, but NAs monotherapy showed a trend towards longer times. During follow-up, patients on NAs monotherapy had a significantly lower rate of HBsAg seroreversion compared to IFN monotherapy patients (4.9% vs. 33.3%).

The one-year and three-year probabilities of confirmed HBsAg seroclearance were significantly higher with NAs therapy (one-year: 0.960 vs. 0.691; three-year: 0.931 vs. 0.641), indicating a more sustained functional cure. NAs monotherapy was independently associated with a lower risk of HBsAg seroreversion.

Most patients experienced HBsAg seroreversion within the first year after seroclearance. IFN monotherapy patients with positive anti-HBs had a higher probability of confirmed HBsAg seroclearance during follow-up, while this difference was not seen in NAs monotherapy patients.

This study provides valuable insights into the durability of HBsAg



seroclearance induced by NAs or IFN monotherapy, with NAs monotherapy showing better outcomes and lower risk of recurrence. Age and ALT levels were also identified as factors associated with HBsAg seroreversion. Understanding these findings can guide clinicians in making informed treatment decisions for patients with chronic Hepatitis B, ensuring better clinical outcomes.

The researchers anticipate that these findings will contribute to the development of more effective and personalized therapeutic strategies for CHB <u>patients</u> in the future. The study highlights the importance of continued research in this area and the potential for leveraging electronic health record data to gain deeper insights into disease management and treatment outcomes.

More information: Zongqi Shi et al, Durability of Hepatitis B surface antigen seroclearance in patients experienced nucleoside analogs or interferon monotherapy: A real-world data from Electronic Health Record, *Genes & Diseases* (2023). DOI: 10.1016/j.gendis.2022.03.003. www.sciencedirect.com/science/ ... 2352304222000563?via%3Dihub

Provided by Chongqing Medical University

Citation: Exploring durability of hepatitis B surface antigen seroclearance using real-world data from electronic health records (2023, September 19) retrieved 12 May 2024 from <u>https://medicalxpress.com/news/2023-09-exploring-durability-hepatitis-surface-antigen.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.