


High levels of particulate air pollution associated with increased breast cancer incidence

September 11 2023



Restricted cubic spline analysis of the association between $PM_{2.5}$ concentrations (1980-1984) and incidence of breast cancer in the NIH-AARP Diet and Health Study. Credit: *JNCI: Journal of the National Cancer Institute* (2023). DOI: 10.1093/jnci/djad170

Researchers at the National Institutes of Health found that living in an area with high levels of particulate air pollution was associated with an increased incidence of breast cancer. The study, published in *JNCI: Journal of the National Cancer Institute*, is one of the largest studies to date looking at the relationship between outdoor air pollution, specifically fine particulate matter, and breast cancer incidence.

The research was done by scientists at the National Institute of Environmental Health Sciences (NIEHS) and the National Cancer Institute (NCI), both part of NIH.

The researchers saw that the largest increases in breast cancer incidence was among women who on average had higher particulate matter levels ($PM_{2.5}$) near their home prior to enrolling in the study, compared to those who lived in areas with lower levels of $PM_{2.5}$.

Particulate matter is a mixture of solid particles and liquid droplets found in the air. It comes from numerous sources, such as motor vehicle exhaust, combustion processes (e.g., oil, coal), wood smoke/vegetation burning, and industrial emissions. The particulate matter pollution measured in this study was 2.5 microns in diameter or smaller ($PM_{2.5}$), meaning the particles are small enough to be inhaled deep into the lungs.

The Environmental Protection Agency has a website known as [Air Now](#)

where residents can enter their zip code and get the air quality information, including PM_{2.5} levels, for their area.

"We observed an 8% increase in breast cancer incidence for living in areas with higher PM_{2.5} exposure. Although this is a relatively modest increase, these findings are significant given that air pollution is a ubiquitous exposure that impacts almost everyone," said Alexandra White, Ph.D., lead author and head of the Environment and Cancer Epidemiology Group at NIEHS. "These findings add to a growing body of literature suggesting that air pollution is related to breast cancer."

The study was conducted using information from the NIH-AARP Diet and Health Study, which enrolled more than 500,000 men and women between 1995–96 in six states (California, Florida, Pennsylvania, New Jersey, North Carolina, and Louisiana) and in two metropolitan areas (Atlanta and Detroit). The women in the cohort were on average about 62 years of age and most identified as being non-Hispanic white. They were followed for approximately 20 years, during which 15,870 breast cancer cases were identified.

The researchers estimated annual average historical PM_{2.5} concentrations for each participant's residence. They were particularly interested in air pollution exposures during a period of 10–15 years prior to enrollment in the study, given the length of time it takes for some cancers to develop. Most previous studies have assessed breast cancer risk in relation to air pollution around the time of study enrollment and did not consider past exposures.

"The ability to consider historic air pollution levels is an important strength of this research," said Rena Jones, Ph.D., senior author and principal investigator of the study at NCI. "It can take many years for breast cancer to develop and, in the past, air pollution levels tended to be higher, which may make previous exposure levels particularly relevant

for cancer development."

To consider how the relationship between air pollution and breast cancer varied by the type of tumor, the researchers evaluated estrogen receptor-positive (ER+) and -negative (ER-) tumors separately. They found that PM_{2.5} was associated with a higher incidence of ER+ breast cancer, but not ER-, tumors. This suggests that PM_{2.5} may affect breast cancer through an underlying biologic pathway of endocrine disruption. ER+ tumors are the most common tumors diagnosed among women in the United States.

The authors note that the study was limited in its ability to explore any differences in the relationship between air pollution and breast cancer across the different study areas. They suggest future work should explore how the regional differences in air pollution, including the various types of PM_{2.5} women that women are exposed to, could impact a woman's risk of developing [breast cancer](#).

More information: Alexandra J White et al, Ambient fine particulate matter and breast cancer incidence in a large prospective US cohort, *JNCI: Journal of the National Cancer Institute* (2023). [DOI: 10.1093/jnci/djad170](#)

Provided by National Institutes of Health

Citation: High levels of particulate air pollution associated with increased breast cancer incidence (2023, September 11) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-09-high-particulate-air-pollution-breast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.