

Neural activity associated with motor commands changes depending on context

September 30 2023



Credit: CC0 Public Domain

Standing at a crosswalk, the signal changes from "don't walk," to "walk." You might step out into the street straight away, or you might look both ways before you cross.

In either scenario, you see the light change, you cross the street. But the <u>context</u> is different; in one case, you didn't think twice. In the other, you waited; looked to the left and right; saw the coast was clear; then stepped



into the street.

Researchers have known that certain brain activity when you see the light change and certain <u>brain activity</u> when you step out into the street are the same no matter the context—there's a known "pathway" that a neuron's activity travels.

Neeraj Gandhi, a bioengineering professor in the Swanson school of Engineering, and team wanted to know, does anything happen along that pathway between the time you see the light change—a stimulus—and the moment you step into the street—an action? Or does the pathway for "crossing the street" look the same, no matter the context?

"If there are two different contexts, even though you're making exactly the same movement, the <u>neural activity</u> in the brain is different." Gandhi said. "In addition to the motor/action command, there is other activity there that tells you something about what's going on cognitively in a given structure," he said.

The findings <u>were published</u> September 29 in the *Proceedings of the National Academy of Sciences (PNAS)*.

From an engineering standpoint, Gandhi said, the finding may have implications for algorithm design. For instance, a similarly designed system could serve as a framework for an autonomous vehicle system that could accelerate when a light turns green, but also delay that action if it sensed something in the crosswalk. The system could analyze the object and, if the coast is clear, it could then begin to drive.

Lead author Eve Ayar, a Ph.D. student at Carnegie Mellon University and a member of Gandhi's lab, found their results may have implications for better understanding the mechanisms underlying <u>executive functions</u> —and ways in which it may be impaired.



"There are a lot of disorders out there where people are unable to take in that sensory stimulus in your environment and make some kind of movement or action in response to that," Ayar said. Soon we may be able to build models that help us better understand how these systems work, and the ways in which they can be disrupted.

"I think this is valuable not only for better understanding this structure of the brain, but potentially it will help us understand how other regions in the <u>brain</u> are operating as well," Ayar said. "And help us be able to differentiate different signals underlying different behaviors."

More information: Eve C. Ayar et al, Distinct context- and contentdependent population codes in superior colliculus during sensation and action, *Proceedings of the National Academy of Sciences* (2023). <u>DOI:</u> <u>10.1073/pnas.2303523120</u>

Provided by University of Pittsburgh

Citation: Neural activity associated with motor commands changes depending on context (2023, September 30) retrieved 13 May 2024 from <u>https://medicalxpress.com/news/2023-09-neural-motor-context.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.