

Finding the balance: Opioids and pain control after surgery

September 11 2023, by Vincent Jacobbi



Credit: Pixabay/CC0 Public Domain

In a recent Mayo Clinic [study](#) published in the *Annals of Surgery*, researchers found that most patients prescribed fewer opioids after surgery were able to maintain satisfactory comfort levels without

requiring more prescription refills later.

Under new evidence-based guidelines, patients undergoing various surgeries received fewer [opioid](#) pills, and 88% reported feeling "very" or "somewhat" satisfied with their [pain management](#).

Researchers worked with the Mayo Clinic Survey Research Center to survey patients undergoing a range of elective surgeries. The survey investigated their post-surgery experiences, overall [pain](#) management approach and opioid use.

The study's results were used to understand the connection between reducing opioid prescriptions and patient satisfaction with [pain relief](#).

Guideline development

The [guidelines were developed](#) through research that examined the amount of opioid medication used by patients, the number of pills left over and the frequency of refills. The results were then applied to create recommendations for prescribing opioids after surgery.

"We wanted to better understand how many pills each patient needed and how many were left over, which set the foundation for publishing guidelines to say how much people need to be prescribed," says senior author of the study Cornelius A. Thiels, D.O., a surgical oncologist at Mayo Clinic.

The guidelines recommend either low, standard or high dosing for each procedure based on specific patient factors associated with variations in opioid requirements.

"We implemented these evidence-based guidelines into practice and developed [educational content](#) for pharmacists, surgeons, [nurse](#)

[practitioners](#), physician's assistants and residents, and saw excellent compliance with the guidelines," says Dr. Thiels.

Impact on pain control

The study showed that while most patients experienced reasonable pain control with reduced opioid prescribing guidelines, a small group of patients did not find their needs met. That group may benefit from further tailoring and individualizing the opioid prescribing guidelines, the study concluded.

While the patient-focused survey supported managing pain with fewer opioids, it also underscored the need for accurate prescribing from the start. The researchers noticed that despite education efforts encouraging patients to dispose of unused opioids properly, many still had leftover pills, which could lead to misuse.

Misuse of prescription opioids remains an important contributor to the opioid epidemic in the U.S., with the study noting that in 2020 alone, 2.3 million people were reported to have a prescription opioid use disorder. That same year, over 16,000 people died from an overdose of prescription opioids.

Dr. Thiels says the findings underscore the need to balance managing pain and [opioid use](#) and the importance of educating patients and prescribing the right amount of medication in the first place.

Hallbera Gudmundsdottir, M.D., first author of the study, adds that the research "is an important first step toward not only safer but also better pain management for every patient after surgery."

Dr. Thiels and Dr. Gudmundsdottir note that future research will focus on using [artificial intelligence](#) to help achieve optimal pain control for

every patient after discharge.

More information: Hallbera Gudmundsdottir et al, Finding the Balance Between Reduced Opioid Prescribing and Patient-reported Pain Management Among General Surgery Patients, *Annals of Surgery* (2023). [DOI: 10.1097/SLA.0000000000005680](https://doi.org/10.1097/SLA.0000000000005680)

Provided by Mayo Clinic

Citation: Finding the balance: Opioids and pain control after surgery (2023, September 11) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-09-opioids-pain-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.