


# Researchers develop novel photo-oxidation therapy for anticancer treatment

September 20 2023



The CityU team developed a new class of near-infrared-activated platinum(IV) photo-oxidants that can effectively eliminate cancer cells in an oxygen-independent manner. The photo-oxidants trigger nonclassical necrosis, which can overcome the resistance of cancer cells to conventional photodynamic therapies and chemotherapies, and activate the immune system. The research highlights the importance of the photo-oxidation capability of metal-based anticancer agents and provides a new direction for the development of next-generation anti-cancer drugs. Credit: Deng, Z. et al., source: <https://www.nature.com/articles/s41557-023-01242-w>

A research team led by scientists from City University of Hong Kong (CityU) has achieved a significant breakthrough by inventing a new class of near-infrared-activated photo-oxidants that can effectively kill cancer cells without requiring oxygen. The photo-oxidants induce a unique form of cancer cell death that can overcome cancer cell resistance. The findings offer a new strategy, called photo-oxidation therapy, and provide a promising direction for the development of anti-cancer drugs.

The paper, titled "Near-infrared-activated anticancer platinum(IV) complexes directly photo-oxidize biomolecules in an [oxygen](#)-independent manner," was published in the journal [Nature Chemistry](#).

Photodynamic therapy, an innovative cancer treatment approach, utilizes photosensitizers to generate [reactive oxygen species](#) (ROSs), which when irradiated by light, selectively kill cancer cells. However, most existing photodynamic therapies rely on the presence of oxygen, while solid cancer tumors often feature a hypoxic microenvironment with very low oxygen levels, limiting the therapeutic efficiency of this approach.

To address this limitation, a research team led by Professor Zhu Guangyu, in the Department of Chemistry, and Professor He Mingliang, in the Department of Biomedical Sciences (BMS) at CityU, discovered an effect called metal-enhanced photo-oxidation. By conjugating metals like platinum with organic photosensitive ligands, they significantly enhanced the photo-oxidation capability. This breakthrough led them to develop a new class of near-infrared-activated platinum(IV) photo-oxidants (Pt(IV) photo-oxidants) that can be activated by near-infrared (NIR) light to directly oxidize biomolecules and effectively kill cancer cells without the need for oxygen.

In their experiments, the team administered Pt(IV) photo-oxidants to mice with tumors through intravenous injection. Four hours later, they applied near-infrared radiation to the mice to activate the photo-oxidants

to attack the cancer cells. The results demonstrated a significant reduction in tumor volume and weight of 89% and 76%, respectively, indicating the potent tumor-inhibitory effect of the Pt(IV) photo-oxidants.

"Intriguingly, we found that the 'death mode' of cancer cells induced by the Pt(IV) photo-oxidants differs from that of any other anticancer agents," said Professor Zhu. "A unique mode of cancer cell destruction was initiated through the dual-action effect of strong intracellular oxidative stress and reduced intracellular pH value."

Their experimental data show that after the Pt(IV) photo-oxidants that accumulated in the [endoplasmic reticulum](#) inside the cancer cells were activated by near-infrared radiation, they vigorously oxidized crucial biomolecules inside the cancer cells without requiring oxygen, generating ROSs, lipid peroxides and protons. The ROSs and lipid peroxides then triggered intensive oxidative bursts, while the protons lowered the intracellular pH value, creating an unfavorable acidic microenvironment for the cancer cells.

In addition, their experiments confirmed that Pt(IV) photo-oxidants effectively activate the [immune system](#) in both in vitro and in vivo settings. The Pt(IV) photo-oxidants triggered immunogenic cell death, stimulating the proliferation and activation of immune cells. The number of T helper and T killer cells, which are crucial for triggering the body's immune response, in the mice treated with photoactivated Pt(IV) photo-oxidants increased by 7- and 23-fold, respectively, compared to the control group.

"By inducing nonclassical necrosis, Pt(IV) photo-oxidants can overcome the resistance of [cancer](#) cells to traditional photodynamic therapies and chemotherapy agents, activate the immune system, and effectively eliminate [cancer cells](#)," explained Professor Zhu.

"These findings serve as proof of concept and suggest that the development of photo-oxidants based on metal-enhanced photo-oxidation is a promising new direction for developing metal-based [anticancer drugs](#)," said Professor He.

The research team plans to conduct preclinical studies to fully characterize the chemical, biological and pharmaceutical properties of the newly invented Pt(IV) photo-oxidants. Their ultimate goal is to identify [lead compounds](#) for clinical testing.

**More information:** Zhiqin Deng et al, Near-infrared-activated anticancer platinum(IV) complexes directly photooxidize biomolecules in an oxygen-independent manner, *Nature Chemistry* (2023). [DOI: 10.1038/s41557-023-01242-w](#)

Provided by City University of Hong Kong

Citation: Researchers develop novel photo-oxidation therapy for anticancer treatment (2023, September 20) retrieved 28 April 2024 from <https://medicalxpress.com/news/2023-09-photo-oxidation-therapy-anticancer-treatment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.