

Q&A: Researcher discusses adverse childhood events and risk of head or neck injury and concussion

September 7 2023



Credit: Unsplash/CC0 Public Domain

Altaf Saadi, MD, MSc, principal investigator of the Neurodisparities &

Health Justice Lab in the Department of Neurology at Mass General and an assistant professor of Neurology at Harvard Medical School, is lead author of a new study in the *Journal of Head Trauma and Rehabilitation*, titled "Examining the Association Between Adverse Childhood Experiences and Lifetime History of Head or Neck Injury or Concussion in Children From the United States." In this Q&A, he discusses the findings.

What question were you investigating?

Is there an association between [adverse childhood experiences](#) (ACEs) and a history of early childhood mild [head](#) or neck injury and concussion?

What methods or techniques did you use to investigate?

We looked at a nationally representative cohort of data from 11,230 children (ages 9 and 10) who are enrolled in the Adolescent Brain Cognitive Development (ABCD) Study.

ACEs are traumatic or stressful events experienced during the first 18 years of life that can have negative physical and mental health consequences later in life.

For this study, we measured eight ACEs:

- Sexual abuse
- Physical abuse
- Emotional neglect
- Parent domestic violence
- Parent substance use disorder

- Parent [mental illness](#)
- Parent criminal involvement
- Parent divorce

The primary outcomes were:

- Head or neck injury
- Concussion

What were your findings?

Having a higher overall ACE count was associated with higher odds of experiencing a head or neck injury and concussion, and that likelihood increased when more ACEs were reported.

- Children with two ACEs had 24% greater odds of head or neck injury and 64% greater odds of concussion
- Children with four or more ACEs had 70% greater odds of head or neck injury and 140% greater odds of concussion

The individual ACE categories of [sexual abuse](#), parent [domestic violence](#), parent mental illness and parent criminal involvement were significantly associated with increased risk of head or neck injury, and parent mental illness with increased risk of concussion.

What are the clinical implications?

ACEs are associated with early childhood head and neck injury or mild concussion and should be integrated into head injury prevention and intervention efforts.

ACEs are associated with many adverse health and psychosocial

outcomes during childhood, adolescence and adulthood. This study adds one more negative outcome to the list (increased risk of head and neck injury or concussion) and further emphasizes the need for more efforts to reduce the prevalence of ACEs.

More information: Altaf Saadi et al, Examining the Association Between Adverse Childhood Experiences and Lifetime History of Head or Neck Injury and Concussion in Children From the United States, *Journal of Head Trauma Rehabilitation* (2023). [DOI: 10.1097/HTR.0000000000000883](https://doi.org/10.1097/HTR.0000000000000883)

Provided by Massachusetts General Hospital

Citation: Q&A: Researcher discusses adverse childhood events and risk of head or neck injury and concussion (2023, September 7) retrieved 12 May 2024 from <https://medicalxpress.com/news/2023-09-qa-discusses-adverse-childhood-events.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
