


Signs of antimicrobial resistance detected in wastewater of care and retirement homes

October 4 2023



Distribution and percentage of sequenced *E. coli* isolate MLSTs per facility and retirement village. Facility 1, n = 24; Facility 2, n = 7; and Retirement, n = 4. Credit: *Microbiology Spectrum* (2023). DOI: 10.1128/spectrum.00731-23

A [new study](#) analyzing wastewater samples from several aged care and retirement homes in Adelaide, has uncovered worrying signs of antimicrobial resistance (AMR) in at least one facility. The study has been published in *Microbiology Spectrum*.

High levels of bacterial resistance against three [common antibiotics](#)—ceftazidime, cefepime and ciprofloxacin—were identified in one aged care residential home. A second facility recorded above average levels of antimicrobial resistance to gentamicin, putting residents' health at risk.

The listed antibiotics are used to treat a variety of bacterial infections, including pneumonia, gynecological, urinary and respiratory tract infections, and those affecting bones and joints.

University of South Australia microbiologist, Associate Professor Rietie Venter, who led the study, says AMR is a concerning trend in aged care facilities.

"Antimicrobial resistance is projected to lead to 300 million deaths worldwide by 2060, and aged care residents are among the most vulnerable due to frequent, inappropriate use of medicines," Assoc Prof Venter says.

Although the wastewater study was confined to three sites and 300 residents, the findings suggest a much wider problem, and are a clear warning to aged care facilities to implement stricter policies when it comes to medication use.

"As well as increasing [death rates](#), AMR can lengthen illness recovery times, especially for immunocompromised people who make up a high proportion of people in aged care homes."

Accurately monitoring the misuse and overuse of antibiotics in residential aged care homes is challenging, hence the use of wastewater-based surveillance, believed to be a first for this sector.

According to the [Australian Commission on Safety and Quality in Health Care](#), general rates of antimicrobial resistance have not

significantly changed in the past two years. However, there are exceptions to this, including increasing resistance to ceftriaxone and fluoroquinolones. Both antibiotics are widely used in aged care settings, despite clinical guidelines recommending them as 'last resort' drugs.

"The results of this study highlight the need for ongoing surveillance of residential [aged care](#) facilities when it comes to medication use," Assoc Prof Venter says. "Given our aging population, there is a crucial need to regularly monitor these facilities and mitigate the threat of AMR."

More information: Sylvia A. Sapula et al, The scope of antimicrobial resistance in residential aged care facilities determined through analysis of *Escherichia coli* and the total wastewater resistome, *Microbiology Spectrum* (2023). [DOI: 10.1128/spectrum.00731-23](https://doi.org/10.1128/spectrum.00731-23)

Provided by University of South Australia

Citation: Signs of antimicrobial resistance detected in wastewater of care and retirement homes (2023, October 4) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-10-antimicrobial-resistance-wastewater-homes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--