


Clinical trial reveals benefits of inhaled nitric oxide for patients with respiratory failure due to COVID-19 pneumonia

October 6 2023, by Noah Brown



This figure depicts the daily cumulative incidence of hypoxemia-free patients from the time of randomization, measured as daily $\text{PaO}_2/\text{FiO}_2$ ratio above 300 mmHg for at least 24 hours. The treatment group (N=94) and the control group (N=99) have been depicted in red and blue, respectively. The data are represented as fractions of 1. Credit: *American Journal of Respiratory and Critical Care Medicine* (2023). DOI: 10.1164/rccm.202304-0637OC

Inhaled nitric oxide gas widens blood vessels in the lungs and is used to treat severe cardio-pulmonary conditions in newborns and adults.

A recent multicenter international phase II clinical trial led by investigators at Massachusetts General Hospital (MGH) and the University of Alabama in Birmingham (UAB) has revealed that inhaled nitric oxide can benefit patients with [respiratory failure](#) due to COVID-19 pneumonia.

The investigator-initiated multi-center trial, which commenced recruitment in March 2020, stands as a testament to the collaborative spirit that prevailed even at the height of the pandemic. The [findings](#) are published in the *American Journal of Respiratory and Critical Care Medicine*.

Prior clinical trials have shown that low-dose (1–20 parts per million) inhaled nitric oxide boosts [blood oxygen levels](#) in [critically ill patients](#) with [acute lung injury](#).

Low-dose inhaled nitric oxide also improved oxygenation during the [severe acute respiratory syndrome](#) (SARS) outbreak of 2003, and recent laboratory experiments showed that nitric oxide inhibits replication of SARS-CoV-2.

To test the potential of inhaled nitric oxide gas to treat SARS-CoV-2–infected patients who develop lung complications, investigators randomized 193 adults with COVID-19 and respiratory failure who were on ventilators at four hospitals in the United States and one in Sweden to receive [high-dose](#) (up to 80 parts per million) nitric oxide for 48 hours or to receive usual care.

Patients who received nitric oxide experienced improved oxygenation and had fewer risks of long-term sensory and motor neurologic symptoms (such as weakness or pain in the extremities) compared with patients receiving usual care.

They also had lower levels of SARS-CoV-2 in their blood and sputum, suggesting that nitric oxide has anti-viral activity. Inhaled nitric oxide was well tolerated and was not linked to any serious adverse events.

Despite these findings, the duration of ventilation and the rates of death at 28 and 90 days did not differ significantly between patients on nitric oxide and those receiving usual care.

"In addition to the well-documented effects on oxygenation, these findings suggest new and promising advantages of utilizing high-concentration inhaled nitric oxide as an antimicrobial agent," says co-senior author Lorenzo Berra, MD, who is the medical director of Respiratory Care at MGH and the Reginald Jenney Associate Professor of Anaesthesia at Harvard Medical School.

"These results highlight the importance of planning future dose-response investigations into the anti-microbial and clinical properties of high-dose inhaled nitric oxide therapy in adults with acute hypoxemic respiratory failure."

"The outcomes of this trial will serve as a critical reference for future research, shedding light on specific subgroups like Black and obese individuals, in whom high-dose inhaled nitric [oxide](#) may yield more substantial clinical benefits," said co-senior author Pankaj Arora, MD, director of the Cardiovascular Clinical and Translational Research Program and the Cardiovascular and Genetics Clinic Program at the UAB Marnix E. Heersink School of Medicine.

"This is especially relevant given the underlying suppression and relative deficiency of the [nitric oxide](#)-cGMP signaling system in these populations."

The investigator-initiated multi-center trial, which commenced recruitment in March 2020, stands as a testament to the collaborative spirit that prevailed even at the height of the pandemic.

More information: Raffaele Di Fenza et al, High-Dose Inhaled Nitric Oxide in Acute Hypoxemic Respiratory Failure due to COVID-19: A Multicenter Phase 2 Trial, *American Journal of Respiratory and Critical Care Medicine* (2023). [DOI: 10.1164/rccm.202304-0637OC](https://doi.org/10.1164/rccm.202304-0637OC)

Provided by Massachusetts General Hospital

Citation: Clinical trial reveals benefits of inhaled nitric oxide for patients with respiratory failure due to COVID-19 pneumonia (2023, October 6) retrieved 10 May 2024 from <https://medicalxpress.com/news/2023-10-clinical-trial-reveals-benefits-inhaled.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--