

Research finds higher risk of dental caries after bariatric surgery

October 31 2023



Credit: Daniel Frank from Pexels

Those who have had surgical obesity treatment have a higher risk of dental caries than before surgery. They also often experience a general decline in oral health. This has been shown by [a thesis](#) from the

University of Gothenburg.

The aim of the research was to study the oral health of individuals before and after treatment for [severe obesity](#). The participants had a BMI of 40 or more, or 35 or more in combination with other [medical conditions](#), such as diabetes. Treatment involved either surgery or medical treatment.

Medical treatment includes lifestyle advice, dietary treatment, possible drug treatment, and support with increased physical activity. However, surgery has proven more effective for [weight loss](#), and with a growing obesity epidemic in many parts of the world, more and more people are having surgery.

"Around 5,000 cases of obesity surgery are performed each year in Sweden alone, and the trend is increasing," says Negin Taghat, who has defended her [doctoral thesis](#) at the Institute of Odontology at the University of Gothenburg's Sahlgrenska Academy, and who works as a dentist for the Swedish Public Dental Service in Region Västra Götaland. "We were therefore interested in seeing whether there is any change in the oral health of these patients after surgery."

Increased caries risk after surgery

The group of 118 [obese individuals](#) followed in the thesis reveals a pattern whereby higher BMI is associated with higher caries risk according to a rising scale. At the highest BMI values, there was a doubled risk of caries and less regular dental care.

Two years after either surgical or medical treatment, a clear division emerged between the groups. Those who had undergone surgery had gone from an average of 15.0 caries lesions on the surface of the tooth enamel to 19.1. Within the group receiving medical treatment, however,

enamel lesions had decreased.

Another example relates to deeper caries lesions in the dentine, with an average pre-treatment initial value of 4.3 lesions. Two years after treatment, individuals in the surgery group had an average of 6.4 such lesions while those in the [medical treatment](#) group had 4.9.

The associations were statistically significant, even when taking factors such as socioeconomic status and other medical conditions into account.

Important knowledge for health care

"Individuals who have undergone surgical obesity treatment may also experience a variety of oral symptoms and an impact on their oral quality of life," continues Negin. "We saw that almost half of individuals experienced poorer oral health."

Symptoms can include hypersensitive teeth and difficulties with chewing. The situation as a whole can also cause social discomfort.

"Health professionals and dental professionals meet these patient groups in their everyday work. It is extremely important for staff to be aware that [oral health](#) can be affected by both obesity and obesity treatment so that preventive measures can be planned."

The group investigated within the thesis is part of the larger BAriatic [surgery](#) SUBstitution and Nutrition (BASUN) study, initiated by researchers at Sahlgrenska Academy to compare long-term outcomes of medical and surgical obesity treatment.

More information: On oral health before and after obesity treatment.
gupea.ub.gu.se/handle/2077/76802

Provided by University of Gothenburg

Citation: Research finds higher risk of dental caries after bariatric surgery (2023, October 31) retrieved 11 May 2024 from <https://medicalxpress.com/news/2023-10-higher-dental-caries-bariatric-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.