


# Investigating the impact of replacing sedentary time with physical activity on genetic risk of coronary heart disease

October 23 2023



Credit: *Journal of Internal Medicine* (2023). DOI: 10.1111/joim.13715

A research team led by Dr. Youngwon Kim, Assistant Professor of Kinesiology from the School of Public Health at the Li Ka Shing Faculty of Medicine of the University of Hong Kong (HKU), in collaboration with the MRC Epidemiology Unit at the University of Cambridge School of Clinical Medicine, has conducted a large-scale epidemiological study to examine the potential health benefits of substituting sedentary time (e.g., sitting) for physically active time in preventing coronary heart disease across different genetic risk levels.

The study found that replacing [sedentary time](#), such as sitting, with an equivalent amount of physically active time is associated with a lower risk of coronary heart disease, regardless of one's [genetic predisposition](#) to the condition. In particular, individuals with a higher genetic risk of coronary heart disease may experience a more significant reduction in their absolute risk of developing the disease compared to those with a lower genetic risk when substituting physical activity for sedentary time. [These findings have been published in the \*Journal of Internal Medicine\*.](#)

The World Health Organization (WHO) Guidelines underscore the importance of minimizing sedentary behavior for achieving optimal cardiovascular health. While excess sedentary time is recognized as a modifiable risk factor, it remains unclear whether replacing sedentary time with an equivalent amount of physical activity time is associated with lower risk of coronary heart disease across varying levels of genetic risk of coronary heart disease.

This research used extensive genotype and phenotype data including

wrist-worn wearable device data from the UK Biobank study which is an ongoing prospective UK cohort of over 500,000 adults.

The analysis included 77,500 white British participants (57% female) without prevalent cardiovascular events and with valid wrist-worn accelerometry data. Each individual's genetic risk of coronary heart disease was assessed using polygenic risk scores based on 300 genetic variants. Sedentary time, light-intensity physical activity, and moderate-to-vigorous physical activity were quantified using wrist-worn wearable device data.

Three primary findings include:

- Greater wearable-device-measured sedentary time is associated with an increased risk of coronary heart disease, regardless of genetic risk of coronary heart disease.
- Replacing 60 minutes/day of sedentary time with an equivalent amount of moderate-to-vigorous physical activity is associated with about 9% lower relative risk of coronary heart disease. Notably, even replacing as little as one minute/day of sedentary time with the same one minute/day of moderate-to-vigorous physical activity is associated with a less than 1% lower relative risk of coronary heart disease, across all levels of genetic risk.
- Replacing sedentary time with the same amount of moderate-to-vigorous physical activity may confer greater reductions in the absolute risk of developing coronary heart disease events in individuals at high genetic risk than in individuals at low genetic risk.

"Our study provides unique insights into the role that replacing sedentary time (such as sitting time) with an equivalent amount of physical activity time could play in preventing coronary heart disease in individuals of varying levels of genetic risk of coronary heart disease," said Dr.

Youngwon Kim, assistant professor, kinesiology, School of Public Health, HKUMed.

"Our findings corroborate the current WHO recommendations on sedentary behavior and physical activity, but suggest that, compared with individuals at low genetic risk of coronary heart disease, those at high genetic risk may have greater reductions in absolute risk of developing [coronary heart disease](#) by substituting physical activity for sedentary time."

"There are more to reveal on the interaction between genetics and physical activity, and this study sets up a framework through which to further explore it," added Haeyoon Jang, Ph.D. candidate, School of Public Health, HKUMed.

"Our study illuminates how high-risk populations may achieve amplified risk reduction by replacing prolonged sitting with [physical activity](#). These insights could shape future WHO Physical Activity guidelines, highlighting the power of lifestyle interventions in promoting cardiovascular health," stated Qiaoxin Shi, Ph.D. candidate, School of Public Health, HKUMed

"To assess the generalizability of the findings to East Asians, it would be worth repeating the study using large East Asian Biobanks," said Dr. Shiu Lun Au Yeung, assistant professor, epidemiology and biostatistics, School of Public Health, HKUMed.

"Leveraging human genetics can enhance our understanding of the intricate interplay between gene and environment interactions in health outcomes," said Dr. Shan Luo, research assistant professor, epidemiology and biostatistics, School of Public Health, HKUMed.

**More information:** Youngwon Kim et al, Replacing device-measured

sedentary time with physical activity is associated with lower risk of coronary heart disease regardless of genetic risk, *Journal of Internal Medicine* (2023). [DOI: 10.1111/joim.13715](https://doi.org/10.1111/joim.13715)

Provided by The University of Hong Kong

Citation: Investigating the impact of replacing sedentary time with physical activity on genetic risk of coronary heart disease (2023, October 23) retrieved 28 April 2024 from <https://medicalxpress.com/news/2023-10-impact-sedentary-physical-genetic-coronary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.