


# Study finds JAK inhibitors, common treatment for arthritis, are effective

November 1 2023

---



Credit: Unsplash/CC0 Public Domain

According to a new paper in *Rheumatology*, published by Oxford University Press, JAK inhibitors, which doctors have used to treat patients with arthritis despite concerns about the effectiveness of such

drugs, actually do work quite well. In a multicenter, retrospective study Japanese researchers found that the drugs resulted in impressive remission rates in patients, most of whom choose to continue such treatment.

Rheumatoid arthritis is a common autoimmune disease characterized by chronic inflammation of joint linings and results in progressive joint destruction and other systemic complications. The use of biological disease-modifying drugs enables patients to enjoy the achievement of low disease activity and remission. But clinics must administer such drugs through subcutaneous or intravenous routes, which is unpleasant for patients, and over time these drugs commonly become less effective.

Recently scientists have developed Janus kinase (JAK) inhibitors for arthritis treatment. Patients take such drugs orally. Previous research has demonstrated the efficacy and safety of JAK inhibitors in randomized controlled trials.

However, some researchers have questioned the potential efficacy of JAK inhibitors for widespread patient use. In practice, doctors mostly treat patients with JAK inhibitors precisely because those patients have other [health problems](#) and so conventional drugs like methotrexate are less effective on them. Real-world patients have distinctive characteristics compared with the patients recruited in [randomized controlled trials](#).

In the present multicenter, [retrospective study](#), researchers using data from 622 patients treated at seven major university hospitals in Japan compared the efficacy and safety of four common JAK inhibitors: tofacitinib, baricitinib, peficitinib, and upadacitinib.

The researchers here found that approximately one in three patients reached remission, three in four reached at least low disease activity,

with both numbers representing impressive efficacy. They noted that more than 80% of the patients were still on the JAK inhibitor drugs after six months.

They believe that this is particularly relevant given that immunological secondary treatment failure, where drugs cease to be effective because they produce adverse immune system responses in patients, cannot occur with these oral medications. Immunological secondary treatment failure is common in patients who treat their arthritis with drugs like methotrexate.

The paper is titled "Real-world comparative study of the efficacy of Janus kinase inhibitors in [patients](#) with [rheumatoid arthritis](#): the ANSWER cohort study."

**More information:** Shinya Hayashi et al, Real-world comparative study of the efficacy of Janus kinase inhibitors in patients with rheumatoid arthritis: the ANSWER cohort study, *Rheumatology* (2023). [DOI: 10.1093/rheumatology/kead543](https://doi.org/10.1093/rheumatology/kead543)

Provided by Oxford University Press

Citation: Study finds JAK inhibitors, common treatment for arthritis, are effective (2023, November 1) retrieved 13 May 2024 from <https://medicalxpress.com/news/2023-10-jak-inhibitors-common-treatment-arthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---