


# Lactate-producing bacteria inside tumors found to promote resistance to radiation therapy

October 19 2023



Graphical Abstract. Credit: *Cancer Cell* (2023). DOI: 10.1016/j.ccell.2023.09.012

Researchers at The University of Texas MD Anderson Cancer Center have discovered that lactate-producing intratumoral bacteria drives resistance to radiation therapy, suggesting that lactic acid-producing bacteria present in various cancers may serve as novel therapeutic targets.

The study, published today in [Cancer Cell](#), reported that a particular bacterial species, *Lactobacillus iners* (L. iners), caused cancer cells to respond to radiation by rewiring metabolic signaling pathways to resist treatment. The researchers also found that L. iners was associated with poorer clinical outcomes in patients with cervical cancer.

"These lactic acid-producing bacteria are seemingly responsible for changing signaling pathways by priming cancer cells to use lactate instead of glucose to fuel growth and proliferation from oxidative stress following [radiation therapy](#)," said corresponding author Lauren Colbert, M.D., assistant professor of Radiation Oncology.

"This is potentially paradigm shifting, and we currently are working on novel approaches to target these specific intratumoral bacteria. We are hopeful that these efforts will lead us to approaches that can benefit patients across several types of cancer."

Cervical cancer is a human papillomavirus (HPV)-related cancer that originates in the same way and is similar microscopically and genomically to anal and head and [neck cancer](#). These types of cancers

are commonly treated with different amounts of radiation.

Cervical cancers often respond well to treatment, but some patients develop resistance, leading Colbert and her colleagues to examine the role of the tumor microbiome in driving differential responses.

Researchers used microbiome sequencing, targeting [bacterial cultures](#) and in vitro models to study the tumor microbiome in this context.

Cervical cancer offers an opportunity for easier repeated tumor microbiome sampling, allowing the researchers to perform large-scale sequencing analyses and targeted culture of tumor-resident bacteria from 101 patients with [cervical cancer](#) undergoing chemoradiation between September 2015 and March 2022.

They discovered that *L. iners* was associated with poor response to radiation and with decreased recurrence-free survival (RFS) and overall survival (OS), even when accounting for known clinical risk factors, gut microbiome factors and immune environment factors in the tumor and in patients' blood. This led the researchers to also perform targeted culture of this specific bacteria from patient samples.

*L. iners* is part of a group of fermenting bacteria that produce lactic acid as a metabolic end product. In this study, the researchers found that cancer-derived *L. iners* within tumors generate L-lactate, which cancer cells use instead of glucose to fuel metabolic cycles and to combat [oxidative stress](#) that occurs after radiation. Tumor lactate and lactate dehydrogenase (LDH) expression are already associated with aggressive tumor growth and poor survival across many cancer types.

The researchers, co-led by Ann Klopp, M.D., Ph.D., professor of Radiation Oncology, were able to induce treatment resistance by introducing lactate-producing *L. iners* to [cancer cells](#) in vitro, resulting in the rewiring of the tumor cell metabolism to use that increased lactate.

The results also revealed that *L. iners* have functions inside of tumors that are not present in healthy patients, suggesting that some changes in these bacteria themselves may occur prior to cancer development.

While this study focused on *L. iners*, the researchers observed a strong association between other tumor-associated lactic acid-producing bacteria and poor survival across multiple cancers, noting that 40% of species are associated with RFS in subsets of lung, colorectal, skin and head and neck cancers.

Colbert and colleagues currently are developing novel approaches to specifically target these [bacteria](#) inside tumors, including topical applications to deliver anti-cancer drugs and improve patient outcomes for various cancer types.

**More information:** Lauren E. Colbert et al, Tumor-resident *Lactobacillus iners* confer chemoradiation resistance through lactate-induced metabolic rewiring, *Cancer Cell* (2023). [DOI: 10.1016/j.ccell.2023.09.012](#)

Provided by University of Texas M. D. Anderson Cancer Center

Citation: Lactate-producing bacteria inside tumors found to promote resistance to radiation therapy (2023, October 19) retrieved 21 May 2024 from <https://medicalxpress.com/news/2023-10-lactate-producing-bacteria-tumors-resistance-therapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--