

Red meat consumption associated with increased type 2 diabetes risk

October 19 2023



Credit: CC0 Public Domain

People who eat just two servings of red meat per week may have an increased risk of developing type 2 diabetes compared to people who eat fewer servings, and the risk increases with greater consumption,

according to a new study led by researchers from Harvard T.H. Chan School of Public Health. They also found that replacing red meat with healthy plant-based protein sources, such as nuts and legumes, or modest amounts of dairy foods, was associated with reduced risk of type 2 diabetes.

The study is published in the *American Journal of Clinical Nutrition*.

"Our findings strongly support [dietary guidelines](#) that recommend limiting the consumption of red [meat](#), and this applies to both processed and unprocessed red meat," said first author Xiao Gu, postdoctoral research fellow in the Department of Nutrition.

While previous studies have found a link between [red meat consumption](#) and type 2 diabetes risk, this study, which analyzed a large number of type 2 diabetes cases among participants being followed for an extended period of years, adds a greater level of certainty about the association.

Type 2 diabetes rates are increasing rapidly in the U.S. and worldwide. This is concerning not only because the disease is a serious burden, but it also is a major risk factor for cardiovascular and kidney disease, cancer, and dementia.

For this study, the researchers analyzed health data from 216,695 participants from the Nurses' Health Study (NHS), NHS II, and Health Professionals Follow-up Study (HPFS). Diet was assessed with food frequency questionnaires every two to four years, for up to 36 years. During this time, more than 22,000 participants developed type 2 diabetes.

The researchers found that consumption of red meat, including processed and unprocessed red meat, was strongly associated with increased risk of type 2 diabetes. Participants who ate the most red meat

had a 62% higher risk of developing type 2 diabetes compared to those who ate the least. Every additional daily serving of processed red meat was associated with a 46% greater risk of developing type 2 diabetes and every additional daily serving of [unprocessed red meat](#) was associated with a 24% greater risk.

The researchers also estimated the potential effects of substituting one daily serving of red meat for another protein source. They found that substituting a serving of nuts and legumes was associated with a 30% lower risk of type 2 [diabetes](#), and substituting a serving of dairy products was associated with a 22% lower risk.

"Given our findings and previous work by others, a limit of about one serving per week of red meat would be reasonable for people wishing to optimize their [health](#) and well-being," said senior author Walter Willett, professor of epidemiology and nutrition.

In addition to [health benefits](#), swapping [red meat](#) for healthy plant protein sources would help reduce greenhouse gas emissions and climate change, and provide other environmental benefits, according to the researchers.

Other Harvard Chan School authors included Frank Sacks and Frank Hu.

More information: Red meat intake and risk of type 2 diabetes in a prospective cohort study of US females and males, *American Journal of Clinical Nutrition* (2023). [DOI: 10.1016/j.ajcnut.2023.08.021](https://doi.org/10.1016/j.ajcnut.2023.08.021)

Provided by Harvard T.H. Chan School of Public Health

Citation: Red meat consumption associated with increased type 2 diabetes risk (2023, October

19) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-10-red-meat-consumption-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.