


A wave theory for explaining neurochemical balance in the brain

October 30 2023



Acetylcholine wave traveling from lateral (L) to medial (M) in the striatum
Credit: Dr. Joshua Goldberg, Prof. Jeffery Wickens, Dr. Nicolas Tristch


In a new study, a group of researchers led by Dr. Joshua Goldberg from the Hebrew University describe a new kind of neurochemical wave in the brain. Their research, [published](#) in *Nature Communications*, unveils the existence of traveling waves of the neurochemical acetylcholine in the striatum, a region of the brain responsible for motivating actions and habitual behaviors.

The motivation to execute an action is widely thought to depend on the release of the another neurochemical, [dopamine](#), in the [striatum](#). Recent research has shown that dopamine is released in wave-like patterns within the striatum. The team led by Goldberg discovered that acetylcholine is also released in the striatum in wave-like patterns. It has long been thought that in order for the striatum to function properly a balance needs to be maintained between dopamine and acetylcholine release in the striatum, and that the disruption of this balance leads to [movement disorders](#) such as Parkinson's disease.

The new study proposes a mathematical mechanism by which simultaneous waves of acetylcholine and dopamine arise, which may represent how this balance is realized.

The research was conducted using state-of-the-art [genetic tools](#) and advanced imaging techniques, allowing the team to visualize the acetylcholine waves in awake, behaving animals. Additionally, imaging techniques were employed to observe the interaction between acetylcholine and dopamine in vitro.

Through a rigorous mathematical analysis, using reaction-diffusion activator-inhibitor models and [computer simulations](#), the team proposed a model that explains the formation of both acetylcholine (the activator) and dopamine (the inhibitor) traveling waves.



Top: a space (y-axis)–time (x-axis) rendition of the location of the acetylcholine release shows it moving in the mediolateral (ML) direction. Bottom: The velocity changes sign when the direction of the acetylcholine wave is reversed Credit: Dr. Joshua Goldberg, Prof. Jeffery Wickens, Dr. Nicolas Tristch

Key highlights of the study:

- First description of acetylcholine waves: in the striatum of healthy behaving animals.
- Local dopamine release is triggered by individual non-dopamine neurons: The study demonstrated that electrical activation of a single acetylcholine-releasing neuron in the striatum is sufficient to induce local dopamine release in its proximity.
- A novel model for how the two neurochemical waves arise simultaneously: The study proposes a novel mathematical model based on the known interaction between acetylcholine and dopamine in the striatum, that can give rise to the simultaneous generation of these waves.

Finally, the study provides strong testable predictions about the relationship between these two wave phenomena and the neural

mechanism for their formation. The study also proposes that dopamine and acetylcholine axons (which are the very long appendages of the dopamine and [acetylcholine](#) neurons) interact directly and locally in the striatum, which is not how neurons are traditionally thought to interact.

More information: Lior Matityahu et al, Acetylcholine waves and dopamine release in the striatum, *Nature Communications* (2023). [DOI: 10.1038/s41467-023-42311-5](#)

Provided by Hebrew University of Jerusalem

Citation: A wave theory for explaining neurochemical balance in the brain (2023, October 30) retrieved 20 May 2024 from <https://medicalxpress.com/news/2023-10-theory-neurochemical-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--