

Umbilical venous catheterization and bloodstream infection among preterm infants

October 31 2023



Though researchers found the same increasing trend for the incidence of UVCassociated CLABSI when the dwell time increased, as in previous studies, this study showed no significant difference in CLABSI incidence between 0-7 days and 7-15 days of dwell time under the current status of UVC management in China. This might have been due to the high exposure rate of antibiotics among preterm infants in China. Credit: Prof. Mingyan Hei from Beijing Children's Hospital, Capital Medical University



Umbilical venous catheterization (UVC) is vital in neonatal care, as it provides essential intravenous access for preterm infants in neonatal intensive care units (NICUs). However, UVC is also associated with some risks, like catheter-associated infections, with central lineassociated bloodstream infection (CLABSI) being a prominent concern.

The incidence of UVC-associated CLABSI varies widely, and the optimal dwell time for UVC remains uncertain. While some reports suggest that early removal of the catheter followed by placement of a peripherally inserted central-line catheter can mitigate CLABSI risk, the data seems insufficient to draw any conclusions. CLABSI not only poses immediate health threats, but can also lead to long-term consequences, including impaired neurodevelopment in preterm infants, highlighting the need for evidence-based national guidelines.

Recognizing the vital need to mitigate this risk, researchers undertook a comprehensive study to investigate the dwell time and incidence of CLABSI associated with UVC in China. The study, led by Professor Mingyan Hei from Beijing Children's Hospital, Capital Medical University, was conducted from November 2019 to August 2021 and included 2,172 preterm infants with UVC.

"To our knowledge, this is the first national-level study on UV catheter dwell time and CLABSI incidence in China," said Prof. Hei.

This study was published in *Pediatric Investigation*.

Among infants aged 30.0 weeks on average, with a median birth weight of 1,258.5 grams, the median UVC dwell time has been found to be 7 days.

"Though we found the same increasing trend of the incidence of UVCassociated CLABSI when the dwell time increases, as in previous



studies, our study showed no significant difference in CLABSI incidence between 0-7 days and 7-15 days of dwell time under the current status of UVC management in China. This might have been due to the high exposure rate of antibiotics among preterm infants in China," said Prof. Hei.

The study also found that a high rate of antibiotic exposure did not cause any reduction in CLABSI rates, raising questions about the efficacy of antibiotics in preventing these infections. The study's findings underscore the complexity of <u>neonatal care</u>, emphasizing the need for nuanced approaches in antibiotic administration.

Additionally, the research highlighted the top complications associated with UVC, with catheter occlusion and migration of the UVC tip being the most prevalent. These insights are crucial for health care providers, allowing them to anticipate challenges and optimize care strategies for neonatal patients.

One of the study's most significant contributions is its national scope. By encompassing a vast array of NICUs across different provinces, this study provides a comprehensive overview of UVC practices and complications in China. The findings not only benefit the <u>medical</u> <u>community</u> but also serve as a reference for policymakers and health care administrators, guiding the formulation of standardized protocols for UVC procedures in NICUs nationwide.

While this study provides crucial insights, it also raises pertinent questions for future research. As the health care landscape evolves, further studies are warranted to delve deeper into the complexities of neonatal care, including the impact of antibiotics, optimal dwell times, and long-term outcomes for preterm infants undergoing UVC procedures. With its illumination of UVC practices in China and challenge to <u>conventional wisdom</u>, this national study lays the



groundwork for evidence-based improvements in neonatal health care.

More information: Xu Zheng et al, Dwell time and bloodstream infection incidence of umbilical venous catheterization in China, *Pediatric Investigation* (2023). DOI: 10.1002/ped4.12403

Provided by Cactus Communications

Citation: Umbilical venous catheterization and bloodstream infection among preterm infants (2023, October 31) retrieved 14 May 2024 from https://medicalxpress.com/news/2023-10-umbilical-venous-catheterization-bloodstreaminfection.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.