

Age at menopause could determine risk for decline in muscle mass and strength

November 29 2023, by Eileen Petridis



Credit: Pixabay/CC0 Public Domain

A shorter reproductive life span, which is determined by a woman's age

at menarche and menopause, has been associated with a number of adverse health effects, including cardiovascular disease and mortality. A new study suggests that it can also lead to a decline in muscle mass and strength as defined by handgrip strength. Results of the study are published online today in *Menopause*.

Sarcopenia is described as a decline in skeletal muscle mass and function and occurs as part of the aging process. It is projected that by 2045, approximately 72.4% of persons aged older than 65 years will be affected by sarcopenia globally. The presence of sarcopenia is concerning because it can lead to reduced physical capability, decreased quality of life, and cardiorespiratory capacity, as well as an increase in metabolic diseases, falls, disability, and mortality.

Menarche and menopause are both known to affect the female aging process. A previous study found that [postmenopausal women](#) experienced an annual [muscle mass](#) loss of 0.6%. One of the most common ways to screen for sarcopenia is by checking handgrip strength. To date, however, no research is known to have focused on the relationship between handgrip strength and the reproductive period.

In a recent study involving more than 2,300 postmenopausal women aged 45 to 75 years, it was found that the more extended the reproductive period, the lower the risk of low absolute handgrip strength. This trend continued even after controlling for other variables.

Other factors that affected handgrip strength were [household income](#), level of education, length of breastfeeding period, and level of intake of vitamin D and protein. Although the age at menopause was associated with reduced low handgrip strength, no relationship was observed between age at menarche and low handgrip strength.

It's also worth noting that in postmenopausal women, the frequency of

sarcopenia varies with race and age. For example, Asian women tend to experience sarcopenia less often.

Survey results are published in the article "The association between reproductive period and handgrip strength in postmenopausal women: a nationwide cross-sectional study."

"This study showed that a longer reproductive period and later age at [menopause](#) was linked to a lower risk of low handgrip strength in postmenopausal Korean women. This finding may relate to the beneficial effects of estrogen on skeletal muscle. Additional longitudinal studies are needed in different populations to confirm these findings," says Dr. Stephanie Faubion, medical director for The Menopause Society.

More information: Sae Rom Lee et al, The association between reproductive period and handgrip strength in postmenopausal women: a nationwide cross-sectional study, *Menopause* (2023). [DOI: 10.1097/GME.0000000000002283](#)

Provided by The North American Menopause Society

Citation: Age at menopause could determine risk for decline in muscle mass and strength (2023, November 29) retrieved 2 May 2024 from <https://medicalxpress.com/news/2023-11-age-menopause-decline-muscle-mass.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.