


Blood thinner apixaban found to prevent strokes in patients with device-detected atrial fibrillation

November 13 2023



A 12 lead ECG showing atrial fibrillation at approximately 150 beats per minute.
Credit: James Heilman, MD/Wikipedia/CC BY-SA 3.0

The widely available blood thinner apixaban substantially reduced stroke in at-risk patients with a type of atrial fibrillation only detectable by a pacemaker or other implanted cardiac electronic device, a global study has found.

The oral anticoagulant medication, which helps to prevent [dangerous blood clots](#) by thinning the blood, reduced the risk of [stroke](#) and blood clotting by 37% and reduced fatal or disabling stroke by 49% in

individuals with device-detected [atrial fibrillation](#), also known as sub-clinical atrial fibrillation (SCAF). This condition is not easily detectable through standard tests like electrocardiograms, unlike clinical atrial fibrillation.

The study was published on Nov. 12 in the *New England Journal of Medicine* (NEJM) and simultaneously presented at the American Heart Association Scientific Sessions by lead researcher Jeff Healey, a senior scientist at the Population Health Research Institute (PHRI), a joint research institute of McMaster University and Hamilton Health Sciences.

Healey and a global team of researchers conducted the largest and longest-running study of apixaban for people with device-detected atrial fibrillation. The study involved more than 4,000 participants at 288 sites in 16 countries over eight years.

"While we found an increase in [major bleeding](#), this non-fatal bleeding is usually reversible and most patients recover from it," says Healey, co-principal investigator (PI) of the study.

"Considering how much more severe and life-altering the strokes prevented by apixaban are, compared to most cases of bleeding, our findings indicate that apixaban should be considered for patients with stroke risk factors who develop device-detected atrial fibrillation," says Stuart Connolly, a PHRI senior scientist and chair of the study.

Renato Lopes, co-PI of the study, cardiologist, and professor in the Department of Medicine at Duke University School of Medicine says, "Our findings can aid doctors in treating these patients, so they don't suffer disabling and fatal strokes."

This global trial was an academic collaboration between the Duke

Clinical Research Institute and PHRI, leading research institutes in the area of cardiology and clinical trials. In the rapidly evolving landscape of cardiac monitoring, Healey underscores the study's far-reaching impact.

"Our findings also have a broader significance," Healey notes, "as we witness the increasing use of implanted and wearable cardiac monitors, along with consumer devices like the Apple Watch. A future where more at-risk individuals are identified and receive effective stroke prevention treatment is on the horizon."

More information: Jeff S. Healey et al, Apixaban for Stroke Prevention in Subclinical Atrial Fibrillation, *New England Journal of Medicine* (2023). [DOI: 10.1056/NEJMoa2310234](https://doi.org/10.1056/NEJMoa2310234)

Provided by McMaster University

Citation: Blood thinner apixaban found to prevent strokes in patients with device-detected atrial fibrillation (2023, November 13) retrieved 20 May 2024 from <https://medicalxpress.com/news/2023-11-blood-thinner-apixaban-patients-device-detected.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
