


Risk of serious infection even in low-active IBD, finds study

November 1 2023, by Gustav Löfgren



Risk of serious infections in patients with histological inflammation vs histological remission of inflammatory bowel disease (IBD) Credit: *Clinical Gastroenterology and Hepatology* (2023). DOI: 10.1016/j.cgh.2023.10.013

IBD is an umbrella term for chronic inflammatory bowel diseases, with a population prevalence of around 0.5%. The main types of IBD are ulcerative colitis and Crohn's disease. Unlike irritable bowel syndrome (IBS), IBD results in visible damage to the intestinal mucous membrane.

IBD is characterized by intermittent symptoms. Periods of high disease activity are sometimes followed by longer periods of low or no activity.

However, the extent to which IBD patients with low disease activity are also at increased risk of serious infections, including sepsis, has been unclear.

A [new study](#) titled "Histological activity in [inflammatory bowel disease](#) and risk of serious infections: A nationwide study," accepted for publication in the journal *Clinical Gastroenterology and Hepatology*, included data on 55,626 individuals diagnosed with IBD. "Serious infections" referred to infections requiring hospitalization.

The difference between healed and unhealed

The results show that during periods of low disease activity but active gastrointestinal inflammation, known as microscopic inflammation, there was an increased risk of being affected by serious infections compared to periods of microscopically healed intestinal mucosa.

In the case of microscopic inflammation, the number of serious infections was 4.62 per 100 people per year. The corresponding figure for microscopically healed mucosa was 2.53. This corresponded to a 59% relative risk increase for residual microscopic gastrointestinal inflammation, on adjusting the results for various confounders.

Interestingly, the results held true even after adjusting for the prescribed IBD medications, and were otherwise similar regardless of age group, sex, and level of education.

The study's first-named and corresponding author is Karl Mårild, an associate professor in pediatrics at the University of Gothenburg's Sahlgrenska Academy and a consultant at the Gastrointestinal and Liver Clinic at Queen Silvia's Children's Hospital in Gothenburg.

Healing provides important protection

"We have shown that even during periods of microscopic intestinal inflammation, IBD patients have an increased risk of serious infections, including sepsis, compared to periods when they have a microscopically healed mucosa," he explains. "This is also true for patients who appear to have low-active disease in clinical terms, but who have microscopic intestinal [inflammation](#) beneath the surface.

"The results indicate that achieving a fully healed [intestinal mucosa](#) in IBD may reduce the risk of serious infections. This is important, as serious infections currently contribute toward increased morbidity and mortality in both children and adults with IBD."

The results from the study are based on data from a national cohort (ESPRESSO) with information from Swedish health registers, and from the quality register for IBD (SWIBREG) on people in Sweden diagnosed with IBD between 1990 and 2016. This information was linked to data from microscopic intestinal examinations of patients with IBD.

More information: Karl Mårild et al, Histological activity in inflammatory bowel disease and risk of serious infections: A nationwide study, *Clinical Gastroenterology and Hepatology* (2023). [DOI: 10.1016/j.cgh.2023.10.013](#)

Provided by Swedish Research Council

Citation: Risk of serious infection even in low-active IBD, finds study (2023, November 1) retrieved 29 April 2024 from <https://medicalxpress.com/news/2023-11-infection-low-active-ibd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.