

Different pain types in multiple sclerosis can cause difficulty staying active

November 1 2023, by Valerie Goodwin



Credit: CC0 Public Domain

For patients with multiple sclerosis, a regular exercise routine is important for managing symptoms. But due to different causes of chronic pain, physical exercise can be more difficult for some.



Research <u>published</u> in *The Journal of Pain* from the University of Michigan has found that widespread pain with nociplastic features can make engaging in <u>physical activity</u> a painful task for <u>patients</u> with MS.

"Widespread pain with nociplastic features is a chronic and diffuse pain which can be challenging to localize or describe precisely," said Libak Abou, Ph.D., research assistant professor and lead author of the paper.

"In a person with MS, this type of pain arises from altered processing signals within the central nervous system. This is opposed to pain that arises from specific tissue damage, classified as nociceptive pain, or pain related to demyelination and axonal damage, classified as neuropathic pain."

Abou and fellow researchers surveyed patients with MS to see if those with a higher indication of widespread pain with nociplastic features were more likely to be insufficiently active or sedentary when compared to their MS counterparts with no chronic pain, nociceptive pain, or <u>neuropathic pain</u>.

Each of the participants was self-reporting with their data.

The results of the survey showed that those who experienced WPNF in addition to their MS were not sufficiently active due to the <u>chronic pain</u> they were experiencing.

"There is a growing need to consider what type of pain MS patients are experiencing before giving them an exercise plan," said Abou.

"The concept of considering widespread pain with nociplastic features when creating exercise plans for MS is newer but could help many patients get to an activity level that will help ease symptoms without causing them intense pain."



For the future, Abou hopes that clinicians can begin doing screenings for underlying pain mechanisms in patients with MS that are struggling to stay active to help further tailor their physical routines to their personal needs.

"The end goal is to help those with MS maintain their functional independence," said Abou.

"It is also important to remember that these patients will likely need extra support from their physical therapy team to keep them on a path with less <u>pain</u>."

Additional authors include Libak Abou, Daniel Whibley, and Anna L. Kratz from the Department of Physical Medicine and Rehabilitation, Michigan Medicine, University of Michigan, Ann Arbor, Michigan as well as the Institute for Healthcare Policy and Innovation, University of Michigan, Ann Arbor, Michigan; and Daniel J. Clauw from the Department of Anesthesiology, Michigan Medicine, University of Michigan, Ann Arbor, Michigan.

More information: Libak Abou et al, Widespread Pain With Nociplastic Features is an Independent Predictor of Low Physical Activity in People with Multiple Sclerosis, *The Journal of Pain* (2023). DOI: 10.1016/j.jpain.2023.09.005

Provided by University of Michigan

Citation: Different pain types in multiple sclerosis can cause difficulty staying active (2023, November 1) retrieved 10 May 2024 from <u>https://medicalxpress.com/news/2023-11-pain-multiple-sclerosis-difficulty-staying.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.