


Getting to the root of visceral gut pain: Research highlights the role played by glia

November 21 2023



IL-1 β expression is increased in enteric glia after inflammation. Credit: *Science*

Signaling (2023). DOI: 10.1126/scisignal.adg1668

Researchers at Michigan State University may have discovered why visceral pain is so common in people who have experienced inflammation in their guts, including patients with irritable bowel syndrome, or IBS.

Working with mouse models, MSU physiologists showed that [nervous system cells](#) known as [glia](#) can sensitize nearby neurons, causing them to send [pain signals](#) more easily than they did prior to inflammation.

"The glia drop the threshold for activating a neuron," said MSU Research Foundation Professor Brian Gulbransen, whose research team authored the new report in the journal [Science Signaling](#).

"So, something that wasn't painful is now painful," Gulbransen said. "It's like when you put on a shirt after getting a sunburn."

This discovery could help researchers develop therapies to lessen or eliminate visceral [pain](#) by counteracting the glia's sensitizing efforts.

Currently, no medicines on the market are designed to act directly on glia, but [pharmaceutical companies](#) are investigating that approach, Gulbransen said.

The team's work offers new information that could help tap into that potential, but it also comes with an important caveat.

The team didn't exactly measure pain. Rather, what they observed was connected to something called nociception.

Nociception is essentially the signals the [nervous system](#) sends in response to physical stimulus. Pain is related but also includes how our brains interpret those signals.

"In general, nociception is pain, but you can have one without the other," said Gulbransen, who works in the Department of Physiology. "It's a little bit of a funny distinction between the two, but it becomes important when extrapolating results from animals to people."

Still, Gulbransen is excited by his team's finding.

"The major thing here is that it suggests a new mechanism contributing to pain in the gut," he said. "And visceral pain is the most common gastrointestinal issue."

Listening to 'silent' cells

The enteric nervous system, the part of the nervous system running through our [gastrointestinal tract](#), is an important but often overlooked part of our anatomy. In fact, it's been nicknamed our [second brain](#).

So perhaps it's not surprising that "important but often overlooked" also aptly describes glial cells in the gut.

Unlike neurons, glia are not electrically active. As researchers developed and refined techniques to probe neurons, the same methods weren't effective for investigating glia.

"If you tried to record [electrical signals](#) from glia, it was just noise. Nothing interesting happened," Gulbransen said. "Glia were sort of ignored as these silent, passive cells."

It turns out, however, that glia are very active chemically. That is, they

react to many different compounds in the body, and they can release different biochemicals in response.

With the advent of new analytical tools and techniques, researchers have become better equipped to observe these cells. Gulbransen's team took advantage of advances in genetic and chemical techniques to monitor [glial cells](#) before and after inflammation in the gut.

"Normal glia in a healthy gut do not change nerve fiber sensitivity," Gulbransen said. "But inflammation triggers a change."

The team discovered that, when exposed to inflammation, glia began releasing compounds that altered the chemistry of the gut and sensitized nerve fibers.

The investigation was spearheaded by Wilmarie Morales-Soto, who earned her doctorate working on this project in Gulbransen's laboratory. She's now a postdoctoral research fellow at the Mayo Clinic.

The research team also included research associate Jacques Gonzales and William Jackson, a professor in the Department of Pharmacology and Toxicology.

Moving forward, Gulbransen and his team are eager to explore looming questions related to visceral pain.

For example, data show that females are more likely to experience visceral pain than males. Early life adversity, including stress and trauma, can also make people more susceptible to visceral pain.

Glia are implicated in both situations and learning more about them could help innovate new ways to treat gut pain.

More information: Wilmarie Morales-Soto et al, Enteric glia promote visceral hypersensitivity during inflammation through intercellular signaling with gut nociceptors, *Science Signaling* (2023). [DOI: 10.1126/scisignal.adg1668](https://doi.org/10.1126/scisignal.adg1668)

Provided by Michigan State University

Citation: Getting to the root of visceral gut pain: Research highlights the role played by glia (2023, November 21) retrieved 9 May 2024 from <https://medicalxpress.com/news/2023-11-root-visceral-gut-pain-highlights.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|