

## STIM-mediated calcium influx regulates maintenance and selection of germinal center B cells: Study

## November 15 2023



**Graphical Abstract** 

Credit: Journal of Experimental Medicine (2023). DOI: 10.1084/jem.20222178

Germinal centers (GCs) are specialized microenvironments where antigen (Ag)-specific B cells undergo antibody affinity maturation and



clonal expansion. Positive selection of high-affinity GC B cells is driven by Ag internalization through their B cell receptor (BCR) and presentation to follicular helper T (Tfh) cells.

However, the requirements of BCR signaling in GC B <u>cells</u> remain poorly understood. Store-operated  $Ca^{2+}$  entry, mediated by stromal interacting molecule 1 (STIM1) and STIM2, is the main  $Ca^{2+}$  influx pathway triggered by BCR engagement.

Researchers Yutaro Yada, Masanori Matsumoto, Tomohiro Kurosaki, Yoshihiro Baba, and their research group from Osaka University have now shown that STIM-deficient B cells have reduced B cell competitiveness compared to wild-type B cells during GC responses. B cell-specific deletion of STIM proteins decreased the number of highaffinity B cells in the late phase of GC formation.

The research is <u>published</u> in the Journal of Experimental Medicine.

STIM deficiency did not affect GC B cell proliferation and Ag presentation, but led to the enhancement of apoptosis due to the impaired upregulation of anti-apoptotic Bcl2a1. STIM-mediated activation of NFAT was required for the expression of Bcl2a1 after BCR stimulation. These findings suggest that STIM-mediated survival signals after Ag capture regulate the optimal selection and maintenance of GC B cells.

GCs are composed of a dark zone where B cells undergo clonal expansion and somatic hypermutation in the immunoglobulin gene and a light zone where B cells capture Ags on follicular dendritic cells (FDCs) and present these to Tfh cells.

In the light zone, high-affinity GC B cells receive a survival signal through induction of the anti-apoptotic bcl2a1 gene, which is required



for NFAT activation by STIM-mediated Ca<sup>2+</sup> influx upon Ag stimulation. Thus, the B cell survival after Ag recognition in GCs may promote affinity maturation by increasing the probability of receiving Tfh cell assistance.

**More information:** Yutaro Yada et al, STIM-mediated calcium influx regulates maintenance and selection of germinal center B cells, *Journal of Experimental Medicine* (2023). DOI: 10.1084/jem.20222178

Provided by Osaka University

Citation: STIM-mediated calcium influx regulates maintenance and selection of germinal center B cells: Study (2023, November 15) retrieved 12 May 2024 from https://medicalxpress.com/news/2023-11-stim-mediated-calcium-influx-maintenance-germinal.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.