

# Understanding survival factors in critically ill patients on extracorporeal membrane oxygenation

November 16 2023

---



Credit: Unsplash/CC0 Public Domain

Extracorporeal membrane oxygenation (ECMO) is a widely used advanced life support procedure that provides cardiac and respiratory

support to critically ill patients. ECMO use has been increasing exponentially over the last decade as it has shown success in resuscitating patients in critical situations like the COVID-19 pandemic and is now a lifesaving treatment modality in intensive care units (ICUs). However, prolonged ECMO use may be associated with an increased risk of mortality.

Identifying [risk factors](#) for in-hospital [mortality](#) and developing standardized nursing practice guidelines for ECMO management may improve the [survival rates](#) in patients.

In this vein, a group of researchers led by Professor Youn-Jung Son from Chung-Ang University, South Korea, set out to study the prevalence and risk factors associated with the 30-day in-hospital mortality in critically ill adult Korean patients on ECMO. This paper was published in [\*Intensive and Critical Care Nursing\*](#).

"In-hospital mortality due to ECMO support, regardless of its type, has not been well reported in Korea. Thus, it is crucial to provide training to various stakeholders in Korea (e.g., ICU nurses) to enhance collaboration, particularly for the optimal ECMO management to reduce in-hospital mortality or improve the survival rate post- hospital discharge," explains Prof. Son.

They conducted a [retrospective cohort study](#) in a single tertiary-care university hospital between May 2020 and April 2021. Patients were screened using their [electronic medical records](#), and 148 patients aged 18 years or above, who received ECMO support for at least 48 hours, were included in the study.

Based on their 30-day in-hospital survival status following ECMO initiation, the researchers categorized the patients as survivors and non-survivors. Various pre-ECMO and post-ECMO (up to 30 days from the

day of initiation) parameters were collected and compared.

The 30-day mortality rate is used to measure performance in clinical settings. In the current study, the researchers found the 30-day in-hospital mortality rate to be 49.3%, with heart and multi-organ failures identified as the most common causes of death. Furthermore, the average duration of ECMO support was approximately 11 days, and the in-hospital median survival was 13 days. This finding emphasizes the importance of early detection of risk factors for mortality to improve patient outcomes.

Further investigations found significant differences in various pre-ECMO and post-ECMO parameters between survivors and non-survivors. Statistical analyses revealed that ECMO weaning failure, new-onset [renal failure](#), and a lower average mean arterial pressure (MAP) of

Citation: Understanding survival factors in critically ill patients on extracorporeal membrane oxygenation (2023, November 16) retrieved 12 May 2024 from <https://medicalxpress.com/news/2023-11-survival-factors-critically-ill-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--