


Childhood pneumonia is surging—while the germs causing it are known, the effects of coinfections aren't

December 5 2023, by Zania Stamataki



Credit: Pixabay/CC0 Public Domain

You may have heard reports about outbreaks of pneumonia affecting children in <u>the Netherlands</u>, <u>Denmark</u> and parts of <u>the US</u> and <u>China</u>,



which are higher than usual for this time of year. Parents are keeping a watchful eye on the news, while public health officials monitor the global incidence of severe respiratory infections and record cases and causes. However, there is no new virus or other type of new pathogen to worry about in these outbreaks.

Here's what we know.

Pneumonia is inflammation of the lungs, usually caused by bacterial or viral infections. It may affect deeper lung tissues than bronchitis, which is airway tube inflammation, and can have <u>symptoms</u> such as fever, breathing problems, and <u>chest pain</u>, showing up as a white shade in lung chest X-rays. There is no special designation of "<u>white lung pneumonia</u>," <u>scary as it may sound</u>.

This season, known respiratory germs, including <u>respiratory syncytial</u> <u>virus</u> (RSV) and Mycoplasma pneumoniae bacteria, have been recorded and are quite expected for this time of year. Mycoplasma outbreaks flare up <u>every few years</u> and can be <u>resistant</u> to common antibiotics.

Most infected children and adults will develop flu-like symptoms that will clear up on their own after a few days, but some may develop into more serious inflammation of the lungs requiring treatment and, in some cases, hospitalization.

But why do some people get more severe diseases? And are there any differences this season?

Healthy children make the most of passing harmless respiratory viruses to each other at an early age, helping to educate their immune systems and build immunity to similar germs in the future.

Over 200 viruses can cause respiratory infection, including RSV,



adenoviruses, enteroviruses, rhinoviruses, and coronaviruses. Symptoms can be more serious in the young when similar infections have not been previously encountered. It takes about <u>a week</u> for specific immunity to develop its arsenal to a new pathogen.

After the infection is cleared, <u>memory cells</u> remain to protect against future infections. We appreciate that antibodies to common cold viruses such as coronaviruses <u>wane</u> after around two years and RSV-specific antibodies in infants may wane <u>even faster</u>. However, memory cells persist, so to escape immunity, viruses mutate to avoid recognition.

There are no differences in expected viral infections this year, and the known culprits were reported in expected waves based on pre-pandemic levels, according to the World Health Organization. *Mycoplasma pneumoniae* flares are also as expected. However, waning mycoplasma-specific herd immunity has been flagged up with a warning to watch out for the resurgence of severe cases following the pandemic restrictions.

Unknown effects of co-infection

COVID restrictions are a thing of the past and people that mix in schools, nursing homes and other high-risk environments are likely to come across several infectious diseases at the same time.

Studies in children with acute severe hepatitis <u>in the UK</u> and <u>the US</u> showed that multiple <u>viral infections</u> were causing unexpected liver damage. Multiple infections can lead to more severe disease, where even viruses that can be <u>considered harmless</u> can add to tissue injury.

In science labs, it has been shown that being infected with more than one respiratory virus (known as "co-infection") can lead to hybrid viruses. These hybrid viruses may behave differently from the individual viruses—causing different symptoms and perhaps even evading the



immune system.

So what do we need to know to determine if a new dangerous outbreak occurs? The incidence of known infections and symptoms are recorded and reported so that any new, potentially more dangerous variants can be detected. Hospitals can become overwhelmed during <u>flu season</u>, but flu does not circulate in isolation and symptoms can worsen when people are simultaneously infected with more than one respiratory <u>virus</u>.

Respiratory infections can cause lung injury, which takes time to heal. Having one respiratory infection followed by another can prolong this healing time, leading to more severe symptoms and longer illness. This is particularly felt by <u>asthma sufferers</u>, who are advised to take their annual flu jab and avoid catching seasonal infections.

Pneumonia-causing infections are contagious. The COVID pandemic has highlighted tips to avoid catching a respiratory <u>infection</u>, which include avoiding contact with infected people, effective ventilation, masking and hand hygiene. Given what we now know about the impact of "harmless" infections, particularly simultaneous infections with more than one pathogen, it is critical to take action to stop the spread.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Childhood pneumonia is surging—while the germs causing it are known, the effects of co-infections aren't (2023, December 5) retrieved 11 May 2024 from https://medicalxpress.com/news/2023-12-childhood-pneumonia-surgingwhile-germs-effects.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.