

Understanding the neuroendocrine basis for social anxiety–like behavior in male mice





Effects of ER α (left) and ER β (right) knockdown in the LS on two behavioral measurements during the OFT. (A) Cumulative duration of time spent in the center area, and (B) total moving distance. Mice were tested in two consecutive days for 10 minutes each. Credit: *Neuroscience* (2023). DOI: 10.1016/j.neuroscience.2023.11.019



Estradiol (E2), a sex steroid hormone, plays an essential role in social behavior, including regulating social anxiety, which is anxiety experienced when unknown individuals are encountered. In males, testosterone secreted by the testes is converted to E2 in the brain, and the E2 binds to two types of estrogen receptors (ERs), ER α and ER β , to regulate social behavior. However, its neuroendocrine basis has not been understood.

In a new study, <u>published</u> in *Neuroscience*, the role of ER α and ER β expressed in the lateral septum (LS), which regulates social anxiety, was investigated using male mice.

The researchers first investigated the expression of ER α and ER β in LS using genetically modified male mice. ER β -expressing cells in the mice were labeled with red fluorescent protein, which revealed that the distributions of ER α and ER β are different. Furthermore, the researchers investigated the knockdown effects of ER α or ER β gene expression in the LS of male mice during situations of social and nonsocial anxiety.

The results show that <u>social anxiety</u> increases with the inhibition of ER β expression. Additionally, ER α - and ER β -positive cells in the LS projected into different regions of the hypothalamus.

Thus, the researchers concluded that $ER\alpha$ - and $ER\beta$ -expressing cells in LS are distinct cell populations with different localizations and neuronal projections, and the $ER\beta$ population plays a crucial role in <u>neural</u> <u>circuitry</u> that regulates anxiety-like behavior in social situations.

More information: Kansuke Hasunuma et al, Estrogen Receptor β in the Lateral Septum Mediates Estrogen Regulation of Social Anxiety-like Behavior in Male Mice, *Neuroscience* (2023). DOI: 10.1016/j.neuroscience.2023.11.019



Provided by University of Tsukuba

Citation: Understanding the neuroendocrine basis for social anxiety–like behavior in male mice (2023, December 27) retrieved 11 May 2024 from https://medicalxpress.com/news/2023-12-neuroendocrine-basis-social-anxietylike-behavior.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.