


Patients with common form of acute myeloid leukemia report better quality of life when treated with quizartinib

December 11 2023



Acute myelocytic leukemia (AML). Credit: Unsplash/CC0 Public Domain

In a study led by researchers at Sylvester Comprehensive Cancer Center at the University of Miami Miller School of Medicine, patients recently

diagnosed with a common and aggressive form of acute myeloid leukemia reported having improved quality of life when a newly approved drug was part of the treatment plan.

Study results will be released during an [oral presentation](#) at the [65th ASH Annual Meeting and Exposition](#), the American Society of Hematology's conference taking place in San Diego, California, Dec. 9-12. The report has been identified by ASH as a clinically relevant abstract.

Mutations in several genes have been implicated in the development of acute myeloid [leukemia](#), or AML, a type of cancer that starts in the blood-forming cells of bone marrow. About 30% of all cases result from [mutations](#) in a gene called FLT3—FMS-like tyrosine kinase 3—and the majority of these involve "internal tandem duplication," or ITD. The new drug, quizartinib, specifically targets this FLT3-ITD mutation, which is found in about 25% of all AML cases and is associated with poor survival.

"In previous research, we showed that patients treated with quizartinib had improved survival compared with patients who didn't receive the drug. This study further demonstrated that patients treated with quizartinib had better quality of life over time and that the side effects from the drug did not negatively affect the improved quality of life," said Dr. Mikkael A. Sekeres, a Sylvester researcher, chief of the Division of Hematology, and senior author of the abstract presented at ASH. One focus of Sekeres's research is AML in older adults.

Quizartinib was recently approved by the U.S. Food and Drug Administration to be used in combination with chemotherapy for the first and second phases of treatment—called induction and consolidation phases—for adult patients with newly diagnosed FLT3-ITD-positive AML. It also can be used alone as maintenance therapy and has been

approved for similar applications in the European Union and Japan.

The current study, looking specifically at patient-reported quality-of-life outcomes—is a continuation—an exploratory endpoint—of findings coming from QuANTUM-First, a global, Phase 3 clinical trial that evaluated the safety and effectiveness of the drug, marketed as VANFLYTA by Daiichi Sankyo Inc.

"Not only did we find that patients treated with quizartinib lived longer, and they lived better, but those who went on to have a [bone marrow transplant](#) had particularly improved quality of life," said Sekeres, who was a co-author of the [QuANTUM-First findings](#) reported earlier this year in *The Lancet*.

The current abstract's first author, Dr. Esther Natalie Oliva, will present the findings. Oliva is a hematologist and researcher at Grande Ospedale Metropolitano Bianchi Melacrino in Reggio Calabria, Italy.

The authors said QuANTUM-First is the first study to explore the impact of an FLT3-inhibitor as first-line [therapy](#) on patient-reported outcomes. Patients treated with quizartinib who achieve complete remission remain in remission longer, continue treatment longer, and have better quality of life than those who do not receive the drug.

Sekeres said the research team plans to conduct additional studies combining quizartinib with different chemotherapy regimens used to treat patients who have FLT3-positive AML to see if it provides similar improvements in quantity and quality of life.

In addition to senior author Sekeres and first author Oliva, authors included Sudhir Unni, Daiichi Sankyo Inc., Basking Ridge, New Jersey; Francesco Cottone, Ph.D., Daiichi Sankyo Italia, Rome, Italy; Rohan Vashi, Pharm.D., Daiichi Sankyo Inc., Basking Ridge, New Jersey;

Xiaocong Li, Ph.D., OPEN Health Evidence & Access, Bethesda, Maryland; and Dr. Jorge Cortes, Augusta University Medical Center, Georgia.

More information: Patient-Reported Outcomes in Acute Myeloid Leukemia Patients with FLT3-ITD Mutation Receiving Quizartinib Vs. Standard Chemotherapy: Results from the Quantum-First Trial: [ash.confex.com/ash/2023/webpro ... ram/Paper189035.html](https://ash.confex.com/ash/2023/webpro...ram/Paper189035.html)

Provided by University of Miami Leonard M. Miller School of Medicine

Citation: Patients with common form of acute myeloid leukemia report better quality of life when treated with quizartinib (2023, December 11) retrieved 8 May 2024 from <https://medicalxpress.com/news/2023-12-patients-common-acute-myeloid-leukemia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--