

Unraveling the age-related cellular changes in osteosarcoma progression



MMA accumulates in the serum of elderly patients with osteosarcoma and accelerates cancer progression. Credit: *Genes & Diseases* (2023). DOI: 10.1016/j.gendis.2023.02.028

In a <u>study</u> published in the journal *Genes & Diseases*, scientists from Union Hospital, Tongji Medical College, and Huazhong University of



Science and Technology delve into the molecular intricacies of osteosarcoma in the elderly. The team investigated senescent cells, cells that no longer divide or grow, and their influence on tumor progression. A spotlight was cast on methylmalonic acid (MMA), a compound found in elevated levels in the elderly, known to be associated with aggressive cancer characteristics.

The team's <u>exploration</u> aimed to decipher MMA's influence on osteosarcoma, focusing on its interplay with the MAFB gene, observed to be highly active in elderly osteosarcoma specimens. The findings were striking: Elevated MMA levels in the elderly enhanced the growth capabilities of osteosarcoma cells. More crucially, the researchers identified a pioneering molecular pathway in which MMA-induced alterations in the MAFB-NOTCH3 axis expedited osteosarcoma progression through the activation of PI3K-AKT pathways.

When the MAFB-NOTCH3 pathway was inhibited, osteosarcoma treatments using AKT inhibitors became markedly more effective. This work has underscored the significance of MAFB as a novel age-associated biomarker for osteosarcoma, suggesting an innovative therapeutic combination targeting the MAFB-NOTCH3 axis and AKT inhibition.

This research not only bridges our understanding gap about osteosarcoma's distinct progression in <u>elderly patients</u> but also carries the beacon of hope for enhancing treatment efficacy for this age bracket. The journey toward more effective, targeted treatments for elderly <u>osteosarcoma</u> patients has taken a pivotal step forward.

More information: Zhenhao Zhang et al, Cellular senescence-driven transcriptional reprogramming of the MAFB/NOTCH3 axis activates the PI3K/AKT pathway and promotes osteosarcoma progression, *Genes & Diseases* (2023). DOI: 10.1016/j.gendis.2023.02.028



Provided by TranSpread

Citation: Unraveling the age-related cellular changes in osteosarcoma progression (2023, December 4) retrieved 12 May 2024 from <u>https://medicalxpress.com/news/2023-12-unraveling-age-related-cellular-osteosarcoma.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.