


Study: Age-related alterations in the oscillatory dynamics serving verbal working memory processing

January 2 2024



Effect of age on alpha and beta oscillations during the encoding, maintenance, and retrieval phases. Credit: 2023 Springer et al.

A new research paper was published on the cover of *Aging*, titled "[Age-related alterations in the oscillatory dynamics serving verbal working memory processing](#)."

Working memory (WM) is a foundational cognitive function involving the temporary storage of information. Unfortunately, WM is also one of the most sensitive cognitive functions to the detrimental effects of aging. Expanding the field's understanding of age-related WM changes is critical to advancing the development of strategies to mitigate age-related WM declines.

In this new study, researchers Seth D. Springer, Hannah J. Okelberry, Madelyn P. Willett, Hallie J. Johnson, Chloe E. Meehan, Mikki Schantell, Christine M. Embury, Maggie P. Rempe, and Tony W. Wilson from Boys Town National Research Hospital, University of Nebraska Medical Center, Washington University School of Medicine, and Creighton University investigated the neural mechanisms serving WM function in seventy-eight healthy aging adults (range: 20.2–65.2 years) using magnetoencephalography (MEG) and a Sternberg WM task with letter stimuli.

"We hypothesized that [older adults](#) would require stronger engagement of key left hemispheric frontal and parieto-occipital WM hubs. Additionally, we expected that prefrontal activity lateralization (i.e., stronger left hemispheric activity) during WM performance would diminish as a function of age, with older individuals tending to utilize a more bilaterally distributed WM network," the researchers write.

Neural activity during the different phases of the WM task (i.e., encoding, maintenance, and retrieval) were imaged using a time-frequency resolved beamformer and whole-brain statistics were performed. The researchers found stronger increases in theta activity and stronger decreases in alpha and beta activity (i.e., more negative relative

to baseline) as a function of healthy aging.

Specifically, age-related increases in theta activity were detected during the encoding period in the primary visual and left prefrontal cortices. Additionally, alpha and beta oscillations were stronger (i.e., more negative) during both encoding and maintenance in the left prefrontal cortex in older individuals. Finally, alpha and beta oscillations during the retrieval phase were stronger (i.e., more negative) in older participants within the prefrontal, parietal, and temporal cortices.

"Together, these results indicate that healthy aging strongly modulates the neural oscillatory dynamics serving WM function," the authors conclude.

More information: Seth D. Springer et al, Age-related alterations in the oscillatory dynamics serving verbal working memory processing, *Aging* (2023). DOI: [10.18632/aging.205403](https://doi.org/10.18632/aging.205403)

Provided by Impact Journals LLC

Citation: Study: Age-related alterations in the oscillatory dynamics serving verbal working memory processing (2024, January 2) retrieved 8 May 2024 from <https://medicalxpress.com/news/2024-01-age-related-oscillatory-dynamics-memory.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
