


Study finds increased body and abdominal fat cause insulin resistance in teenagers, raising type 2 diabetes risk

January 15 2024



Both total body fat mass and abdominal fat cause hyperglycemia, hyperinsulinemia, and insulin resistance, with abdominal fat doubling the risk of metabolic dysfunction. However, light physical activity can significantly reduce total body fat mass and insulin resistance. Credit: Andrew Agbaje

High fat mass in adolescence causes insulin resistance, which can lead to a vicious cycle of worsening insulin resistance and obesity by young adulthood, a new study shows. However, having a high muscle mass partially protects against insulin resistance.

The study was conducted in collaboration between the Université de Montréal in Canada, the University of Bern in Switzerland, Aarhus University in Denmark, the University of Bristol in the U.K., the University of Exeter in the U.K., and the University of Eastern Finland, and the results were [published](#) in the *Journal of Clinical Endocrinology and Metabolism*.

Childhood and adolescent obesity remains a global epidemic. Estimated by [body mass index](#) (BMI), obesity has been associated with several cardiovascular, neurological, and musculoskeletal diseases as well as type 2 diabetes in adulthood. However, BMI is a poor measure of obesity in childhood and adolescence since it does not distinguish between muscle mass and fat mass.

Insulin resistance occurs when cells in the body fail to take up glucose from the blood in spite of a normal amount of insulin production, leading to overproduction of insulin, known as hyperinsulinemia. Insulin resistance is a precursor of type 2 diabetes already in youth, but [long-term studies](#) relating to the association of total body fat mass, [abdominal fat](#), and muscle mass with the risk of insulin resistance in a large population of young people are lacking.

The current study used data from the University of Bristol's Children of the 90's cohort, also known as the Avon Longitudinal Study of Parents and Children. Altogether 3,160 adolescents—1,546 males and 1,614 females—were included in the analyses. The adolescents were 15 years old at baseline, and they were followed up for nine years until young adulthood at age 24 years. Total body fat mass, abdominal fat, and

muscle mass were measured with dual-energy X-ray absorptiometry at age 15 years, and repeated at age 17 and 24 years. Similarly, fasting glucose and insulin were measured from [blood samples](#) taken at ages 15, 17, and 24 years, and insulin resistance was calculated.

With extensive control for inflammation, lipids, [blood pressure](#), smoking status, sedentary time, [physical activity](#), socio-economic status, and family history of cardiovascular disease, it was observed that each 1 kg cumulative increase in total body fat mass from mid-adolescence through young adulthood increased the risk of excessive blood glucose (hyperglycemia) by 4%, abnormally high insulin level (hyperinsulinemia) by 9%, and insulin resistance by 12%.

Each 1 kg increase in abdominal fat had even more pronounced effects, increasing the risk of hyperglycemia by 7%, hyperinsulinemia by 13%, and insulin resistance by 21%. However, each 1 kg increase in muscle mass reduced the risk of both hyperinsulinemia and insulin resistance by 2%.

It was also found that a high amount of total body fat mass at age 15 years caused high insulin resistance at age 17 years. A high amount of body fat mass at age 17 years caused high insulin resistance at age 24 years, and, simultaneously, high insulin resistance at age 17 years caused high total body fat mass at age 24 years, resulting in a [vicious cycle](#). The results were consistent in both males and females, regardless of BMI.

"This is the first long-term evidence of the morbid danger of high total body and abdominal fat in the young population, with abdominal fat twice as dangerous as total body fat. To observe a vicious cycle of fat mass and insulin resistance between the ages of 17 and 24 years is extremely disheartening. Fat mass contributes 75%, while insulin resistance contributes 25% to the fat mass-[insulin](#) resistance vicious cycle. Therefore, preventing [weight gain](#) is the best way to break this

cycle," says Andrew Agbaje, a physician and pediatric clinical epidemiologist at the University of Eastern Finland.

"Nonetheless, there is good news: We recently established that sedentary time contributes 10% to the total body [fat mass](#) gained in youth years, which can be completely reversed by 3–4 hours/day of light physical activity. Although the increase in [muscle mass](#) lowers [insulin resistance](#) only by a small amount, its protective effect when combined with light exercise may be critical to reducing the risk of type 2 diabetes. This is why teenagers should be encouraged to take up light physical activity as discussed in a recent podcast," Agbaje concludes.

More information: Andrew O Agbaje et al, DXA-based Fat Mass With Risk of Worsening Insulin Resistance in Adolescents: A 9-Year Temporal and Mediation Study, *The Journal of Clinical Endocrinology & Metabolism* (2024). [DOI: 10.1210/clinem/dgae004](https://doi.org/10.1210/clinem/dgae004)

Provided by University of Eastern Finland

Citation: Study finds increased body and abdominal fat cause insulin resistance in teenagers, raising type 2 diabetes risk (2024, January 15) retrieved 27 April 2024 from <https://medicalxpress.com/news/2024-01-body-abdominal-fat-insulin-resistance.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--