

Predicting brace adherence could change the game in scoliosis treatment

January 24 2024, by Joanne Barker



Patients with moderate scoliosis who adhered to bracing in the first month continued to do so 24 months later. Credit: Children's Hospital Boston

When it comes to preventing scoliosis progression, is it possible to make bracing more effective? For decades, spine specialists focused on

improving the braces themselves, making them lighter, less obtrusive, and easier to put on and take off. (The Boston Brace, developed at Boston Children's Hospital in the early 1970s, is one example.)

With the addition of built-in sensors that indicate how many hours a brace was worn, two things became clear. One: Despite improvements in brace design, many patients wore their [braces](#) far less than they thought or said they did. Two: Patients who wore their braces more had better outcomes.

If a [spine](#) specialist could predict which patients were least likely to adhere to a bracing regimen, could they reduce the number who eventually need spinal fusion surgery? Would it be possible, in fact, to increase the rate of curve correction through bracing alone?

Timothy Hresko, MD, director of spine research and [quality improvement](#) in Boston Children's Spine Division, believes the answer to both of these questions is yes.

Predicting who will and who won't wear a brace

As he read reports linking hours in a brace to curve stabilization, Hresko started to rethink the field's approach to moderate scoliosis. Rather than continually trying to improve brace designs, he thought, "maybe it's time to shift our focus to helping kids accept the fact that they need to wear the brace."

But first, spine specialists would need to know more about patterns of adherence. Do patients who initially resist wearing a brace eventually come around? Or does a patient's early acceptance or nonacceptance of bracing indicate how they're likely to behave in the future?

In a [study published](#) in *Spine*, Hresko and his colleagues analyzed brace

adherence in 60 patients with [adolescent idiopathic scoliosis](#) over two years. Patients fell into one of two groups, adherent—defined as wearing the brace for at least 80% of the hours prescribed—and nonadherent.

At their one-month follow-up visit, 32 patients met the criteria of adherence. By the time of their six-month visit, these patients had worn their braces an average of five hours more per day than those in the nonadherent group. This pattern continued throughout the 24 months studied.

Those who started out wearing their braces the most continued to do so while those who resisted wearing a brace early on did not adapt to the treatment regimen over time.



Many patients who adhered to bracing, as this patient did, showed significant curve correction. Credit: Children's Hospital Boston

From scoliosis stabilization to correction

In keeping with previous studies, bracing was far more successful among patients in the adherent group, with only one progressing to severe scoliosis (a Cobb angle greater than 40 degrees). By contrast, seven patients in the nonadherent group progressed from moderate to severe scoliosis. Further, patients in the adherent group showed significant curve reduction.

These results, says Hresko, are reason to change the way spine specialists talk about scoliosis and bracing with patients.

Up until now, patients with moderate scoliosis have typically been told that wearing a brace may prevent their curve from progressing but they shouldn't expect it to cure their scoliosis. By contrast, Hresko presents bracing as an opportunity for patients to take charge of their lives. "It's crucial that patients are actively involved in their own care," he explains. "If they're passive, if someone else is making them wear the brace, it's a lot less likely they'll be adherent."

The future of bracing adherence

Hresko considers adherence a game changer in scoliosis treatment.

"There's a large group of adolescents with scoliosis who can reduce the chance they'll need surgery by spending more hours in their brace," he says.

To understand how to best reach them, Hresko and his team are planning

a prospective study that will analyze factors—familial, social, personal—driving adherence or nonadherence. Ultimately, they plan to use the results to help more [patients](#) with scoliosis reach a place at which wearing a [brace](#) is a choice they make for their own well-being.

More information: Gabriel S. Linden et al, Early Adherence to Prescribed Brace Wear for Adolescent Idiopathic Scoliosis Is Associated With Future Brace Wear, *Spine* (2022). [DOI: 10.1097/BRS.0000000000004446](#)

Provided by Children's Hospital Boston

Citation: Predicting brace adherence could change the game in scoliosis treatment (2024, January 24) retrieved 12 May 2024 from <https://medicalxpress.com/news/2024-01-brace-adherence-game-scoliosis-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--