


# Study finds raised blood pressure is the leading risk factor for death in Australia

February 21 2024

The ranking in the contribution of risk factors towards all-cause death in Australia



The ranking in the contribution of risk factors towards all-cause, CVD, IHD and stroke deaths by males and females in 1990, 2010, and 2019, with percentage change and 95% UI in age-standardised deaths. Credit: Xu et al., 2024, PLOS ONE, CC-BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>)

Raised blood pressure has been the leading risk factor for death in Australia for the past three decades, according to a study published February 21, 2024, in the open-access journal *PLOS ONE* led by Alta Schutte and Xiaoyue Xu from The George Institute for Global Health and UNSW, Sydney, with colleagues across Australia. It is also the main contributor to deaths from cardiovascular disease (CVD) specifically.

Raised blood pressure has long been recognized as a contributing factor to CVD and [death](#), but is not always prioritized in national health plans.

In this study, researchers focused on Australia, which lags behind other high-income countries in hypertension control. Data on how raised blood pressure compares to other [risk factors](#) for CVD burden—and how this changes over time—can help to guide public health agendas and inform the effectiveness of public health policies.

Researchers analyzed epidemiologic data from the Global Burden of Disease (GBD) study between 1990 and 2019 to determine the leading risk factors associated with both all-cause and CVD deaths, over time and between gender and [age groups](#). The GBD study provides data on nearly 400 diseases and 87 risk factors across 204 countries.

They found that while the contribution of raised blood pressure to these outcomes declined early in the study period (from around 54% to around 44%), it persisted as the leading risk factor for all-cause and CVD deaths. Dietary factors and tobacco use rounded out the top three risk factors.

These findings strongly align with the recently established National Hypertension Taskforce of Australia, which aims to improve Australia's blood pressure control rates from 32% to 70% by 2030 (Hypertension—Australian Cardiovascular Alliance ([ozheart.org](http://ozheart.org))). The research findings further advocate for the prioritization of blood

pressure control on the public health agenda.

Differences by gender and age were also seen. For example, the contribution of raised blood pressure to stroke-related deaths in males aged 25–49 years were higher than other age groups, exceeding 60% and increasing steeply over time.

The study reinforces the importance of blood pressure control and awareness. The researchers hope that the data will urge policymakers to prioritize blood pressure control efforts in Australia and will provide insight into age groups and populations that would benefit from more targeted action.

The authors add, "There is no doubt that raised blood pressure has remained the leading risk factor for all-cause and cardiovascular deaths in Australia across the past three decades. Our findings support actions to strengthen [primary care](#) and to improve the prevention, detection, treatment and control of raised [blood pressure](#), with the goal of significantly reducing all-cause and cardiovascular deaths in Australia over the next decade."

**More information:** The contribution of raised blood pressure to all-cause and cardiovascular deaths and disability-adjusted life-years (DALYs) in Australia: Analysis of global burden of disease study from 1990 to 2019, *PLoS ONE* (2024). [DOI: 10.1371/journal.pone.0297229](https://doi.org/10.1371/journal.pone.0297229)

Provided by Public Library of Science

Citation: Study finds raised blood pressure is the leading risk factor for death in Australia (2024, February 21) retrieved 11 May 2024 from <https://medicalxpress.com/news/2024-02-blood-pressure-factor-death-australia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.