

## Largest COVID vaccine study yet evaluates adverse events of special interest

February 19 2024, by Jason Gale, Bloomberg News



Credit: Pixabay/CC0 Public Domain

Vaccines that protect against severe illness, death and lingering long COVID symptoms from a coronavirus infection were linked to small increases in neurological, blood, and heart-related conditions in the



largest global vaccine safety study to date.

The <u>rare events</u>—identified early in the pandemic—included a higher risk of heart-related inflammation from mRNA shots made by Pfizer Inc., BioNTech SE, and Moderna Inc., and an increased risk of a type of blood clot in the brain after immunization with viral-vector vaccines such as the one developed by the University of Oxford and made by AstraZeneca Plc.

The viral-vector jabs were also tied to an increased risk of Guillain-Barre syndrome, a <u>neurological disorder</u> in which the <u>immune system</u> mistakenly attacks the peripheral nervous system.

More than 13.5 billion doses of COVID vaccines have been administered globally over the past three years, saving over 1 million lives in Europe alone. Still, a small proportion of people immunized were injured by the shots, stoking debate about their benefits versus harms.

The new research, by the Global Vaccine Data Network, was <u>published</u> in the journal *Vaccine*, with the data made available via interactive dashboards to show methodology and specific findings.

The research looked for 13 <u>medical conditions</u> that the group considered "adverse events of special interest" among 99 million vaccinated individuals in eight countries, aiming to identify higher-than-expected cases after a COVID shot. The use of aggregated data increased the possibility of identifying rare safety signals that might have been missed when looking only at smaller populations.

Myocarditis, or inflammation of the heart muscle, was consistently identified following a first, second and third dose of mRNA vaccines, the study found. The highest increase in the observed-to-expected ratio



was seen after a second jab with the Moderna shot. A first and fourth dose of the same <u>vaccine</u> was also tied to an increase in pericarditis, or inflammation of the thin sac covering the heart.

## Safety signals

Researchers found a statistically significant increase in cases of Guillain-Barre syndrome within 42 days of an initial Oxford-developed ChAdOx1 or "Vaxzevria" shot that wasn't observed with mRNA vaccines. Based on the background incidence of the condition, 66 cases were expected—but 190 events were observed.

ChAdOx1 was linked to a threefold increase in cerebral venous sinus thrombosis, a type of blood clot in the brain, identified in 69 events, compared with an expected 21. The small risk led to the vaccine's withdrawal or restriction in Denmark and multiple other countries. Myocarditis was also linked to a third dose of ChAdOx1 in some, but not all, populations studied.

Possible safety signals for transverse myelitis—spinal cord inflammation—after viral-vector vaccines were identified in the study. So was acute disseminated encephalomyelitis—inflammation and swelling in the brain and spinal cord—after both viral-vector and mRNA vaccines.

Seven cases of acute disseminated encephalomyelitis after vaccination with the Pfizer-BioNTech vaccine were observed, versus an expectation of two.

The adverse events of special interest were selected based on preestablished associations with immunization, what was already known about immune-related conditions and pre-clinical research. The study didn't monitor for postural orthostatic tachycardia syndrome, or POTS,



that some research has linked with COVID vaccines.

Exercise intolerance, excessive fatigue, numbress and "brain fog" were among common symptoms identified in more than 240 adults experiencing chronic post-vaccination syndrome in a separate study conducted by the Yale School of Medicine. The cause of the syndrome isn't yet known, and it has no diagnostic tests or proven remedies.

The Yale research aims to understand the condition to relieve the suffering of those affected and improve the safety of vaccines, said Harlan Krumholz, a principal investigator of the study, and director of the Yale New Haven Hospital Center for Outcomes Research and Evaluation.

"Both things can be true," Krumholz said in an interview. "They can save millions of lives, and there can be a small number of people who've been adversely affected."

**More information:** K. Faksova et al, COVID-19 vaccines and adverse events of special interest: A multinational Global Vaccine Data Network (GVDN) cohort study of 99 million vaccinated individuals, *Vaccine* (2024). DOI: 10.1016/j.vaccine.2024.01.100

Hannah J. Morgan et al, Acute disseminated encephalomyelitis and transverse myelitis following COVID-19 vaccination—A self-controlled case series analysis, *Vaccine* (2024). DOI: 10.1016/j.vaccine.2024.01.099 , <u>doi.org/10.1016/j.vaccine.2024.01.099</u>

2024 Bloomberg L.P. Distributed by Tribune Content Agency, LLC.

Citation: Largest COVID vaccine study yet evaluates adverse events of special interest (2024, February 19) retrieved 12 May 2024 from <u>https://medicalxpress.com/news/2024-02-largest-</u>



covid-vaccine-adverse-events.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.