

Poorly controlled asthma emits same quantity of greenhouse gas as 124,000 homes each year in the UK: Study

February 27 2024



Credit: Pixabay/CC0 Public Domain

Patients whose asthma is poorly controlled account for eight times excess greenhouse gas emissions compared with those whose condition

is well controlled—equivalent to that produced by 124,000 homes each year in the U.K. This result has been published online in the journal *Thorax*.

Improving the care of asthma patients could achieve substantial carbon emissions savings, and help the NHS meet its net zero target, say the researchers.

Health care is a major contributor to greenhouse gas emissions and in 2020 the NHS set an ambitious target of reducing its [carbon footprint](#) by 80% over the next 15 years, with the aim of reaching net zero by 2045, note the researchers.

Asthma is poorly controlled in around half of those with the condition in the U.K. and Europe, increasing the risk of hospital admission and [severe illness](#) as well as [health care](#) costs.

To gauge the environmental footprint of asthma care in the U.K., the researchers retrospectively analyzed the anonymized health records of 236,506 people with asthma whose data had been submitted to the Clinical Practice Research Datalink between 2008 and 2019.

Greenhouse gas (GHG) emissions, measured as carbon dioxide equivalent (CO₂e), were estimated for asthma-related medication use, health care resource utilization and severe exacerbations during follow-up of patients with asthma.

Well-controlled asthma was categorized as no episodes of severe worsening symptoms and fewer than three prescriptions of short-acting beta-agonists (SABAs) reliever inhalers in a year.

Poorly controlled asthma was categorized as three or more SABA canister prescriptions or one or more episodes of severe worsening

symptoms in a year.

A severe exacerbation of asthma was defined as worsening symptoms requiring a short course of oral corticosteroids, an emergency department visit, or hospitalization.

Excess GHG emissions due to suboptimal asthma control included at least three or more SABA canisters per year, severe exacerbations and any GP visits within 10 days of hospitalization or an emergency department visit.

The researchers calculated that the overall carbon footprint attributed to asthma care when scaled to the entire U.K. asthma population added up to 750,540 metric tons CO₂e/year.

Asthma was poorly controlled in just under half (47%; 111,844) of the patients. And poorly controlled asthma contributed to excess greenhouse gas emissions of 303,874 tons CO₂e/year—equivalent to emissions from more than 124,000 homes in the U.K., they estimate. The excess GHG emissions were eight-fold higher on average for a person with poorly controlled asthma than in the well controlled asthma patients.

The excess GHG emissions were 90% comprised of inappropriate SABA use with the remainder mostly due to health care resource utilization such as GP and hospital visits, required to treat severe worsening symptoms.

Poorly controlled asthma generated three-fold higher greenhouse gas emissions on average for a person with poorly controlled asthma compared with well-controlled asthma when taking into account GHG emissions related to all aspects of asthma care including routine prescribing and management.

The researchers acknowledge various limitations to their findings, including that the study results were largely descriptive in nature. And factors other than the level of asthma symptom control, such as prescribing patterns, may also have contributed to high SABA use.

But they nevertheless write, "Our study indicates that poorly controlled asthma contributes to a large proportion of asthma-care related [greenhouse gas emissions](#) with inappropriate SABA use emerging as the single largest contributor."

The Global Initiative for Asthma no longer recommends SABA used alone as the preferred reliever for acute asthma symptoms, they add.

The authors conclude that efforts to improve [asthma](#) treatment practices including curtailing inappropriate SABA use and implementing evidence-based treatment recommendations, could result in substantial carbon savings.

More information: Greenhouse gas emissions associated with suboptimal asthma care in the UK: the SABINA healthCARE-Based enviroNmental cost of treatment (CARBON) study, *Thorax* (2024). [DOI: 10.1136/thorax-2023-220259](https://doi.org/10.1136/thorax-2023-220259)

Provided by British Medical Journal

Citation: Poorly controlled asthma emits same quantity of greenhouse gas as 124,000 homes each year in the UK: Study (2024, February 27) retrieved 27 April 2024 from <https://medicalxpress.com/news/2024-02-poorly-asthma-emits-quantity-greenhouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.