


Psychedelic experiences linked with improved sexual function

February 7 2024



Percentage of participants who retrospectively rated decreases or increases in sexual interest, arousal, activity, satisfaction, and anxiety (reversed) after treatment with psilocybin or escitalopram at the 6 weeks follow-up of Study 2. Credit: *Scientific Reports* (2024). DOI: 10.1038/s41598-023-49817-4

Magic mushrooms, LSD and other psychoactive compounds may help to

improve sexual function for months after the psychedelic experience.

These are the findings of the first known scientific investigation into the effects of psychedelics on sex, which finds that participants reported improvements across a range of measures for several weeks after an acute psychedelic experience.

In the [study](#), published today in the journal *Scientific Reports*, researchers from the Center for Psychedelic Research at Imperial College London gathered questionnaire responses from almost 300 people before and after a psychedelic experience.

By combining responses from two groups—people taking psychedelics for recreational or wellness/ceremonial purposes, as well as a small group from a clinical trial assessing psilocybin (the compound found in magic mushrooms) for depression—they were able to investigate how the experience influenced a range of aspects relating to sexual functioning.

The analysis reveals that on average, people reported improvements across a range of areas of [sexual function](#) up to six months after their psychedelic experience, including their enjoyment of sex, sexual arousal, satisfaction with sex, attraction to partner, their own physical appearance, communication, and their sense of connection.

In addition, among the small group taking psilocybin in the clinical trial for depression, they found that almost half of participants reported improvements in sexual arousal, interest and satisfaction with sex, while people treated with a leading antidepressant mostly reported decreases in sexual functioning.

According to the team, the findings open the possibility for applications in a range of therapeutic settings—such as couples therapy. They also

highlight further potential benefit in the treatment of depression, where psychedelic compounds such as psilocybin could potentially help to avoid drug-induced [sexual dysfunction](#)—one of the biggest side effects with current gold standard antidepressants.

Tommaso Barba, first author of the study and Ph.D. student based at the Center for Psychedelic Research at Imperial College London, said, "We believe this is the first scientific study to explore the effects of psychedelics on sexual functioning. Our findings suggest potential implications for conditions that negatively affect sexual health, including [clinical depression](#) and anxiety.

"This is particularly significant given that sexual dysfunction, often induced by antidepressants, frequently results in people stopping these medications and subsequently relapsing.

"On the surface, this type of research may seem 'quirky,' but the psychological aspects of sexual function—including how we think about our own bodies, our attraction to our partners, and our ability to connect to people intimately—are all important to psychological well-being in sexually active adults.

"Sexuality is a fundamental human drive. For example, we know that sexual dysfunction is linked to lower well-being in healthy adults, can impact relationship satisfaction, and is even linked to subjective happiness and 'meaning in life.'"

Bruna Giribaldi, senior author on the paper and formerly based at the Center for Psychedelic Research at Imperial, said, "So far, all the research on depression has only focused on whether treatments cause sexual dysfunction. We wanted to make sure we went deeper than that and explored more aspects of sexuality that could be impacted by these treatments.

"We were interested in finding out whether psychedelics could influence people's experiences of sexuality in a positive way, as it appeared from existing anecdotal evidence.

"Sexual side effects are often underreported in clinical trials, as people tend to not directly report them unless explicitly asked. Aware of this, we were determined to thoroughly investigate this area."

In the analysis, researchers combined responses to questionnaires from two studies, with participants providing responses before a psychedelic experience, and then at four weeks and six months after.

In the first of the two studies, researchers recruited people already planning to take psychedelics in the near future, either for recreation or as part of a psychedelic ceremony. Participants provided responses to the online survey before taking substances (including psilocybin/[magic mushrooms](#)/truffles, ayahuasca, DMT, San Pedro, and LSD) and twice after the experience.

A total of 261 people were included in the analysis, and response showed that overall, they were more likely to report improvements across all areas of sexual functions for up to six months after their psychedelic experience. The most significant improvements were in sexual pleasure, satisfaction with their own appearance, satisfaction and communication with their partner, as well as perceiving sex as a "spiritual experience."

The second study, led by the former director of the Center, Professor Robin Carhart-Harris, looked at 59 people taking part in a clinical trial to assess psilocybin for treating major depressive disorder—with 30 people given psilocybin, and 29 taking the antidepressant escitalopram (a type of SSRI).

Participants on the psilocybin arm were more likely to report

improvements across all areas of sexual function at the end of the trial, with the most significant improvements reported in [sexual arousal](#), interest, satisfaction and communication with their partner.

Patients treated with escitalopram tended to report decreases. An exception was satisfaction with their appearance, which improved in both arms. Levels of reported sexual dysfunction were also significantly lower for the psilocybin group (13 % reporting sexual dysfunction at the end of the trial) compared to those in the citalopram group (59% reporting dysfunction).

The authors note that, overall, neither study found a significant change in participants' perceived importance of sex after a psychedelic experience. This indicates that psychedelics don't cause an excessive focus on sex or "hyper-arousal." Instead, they may transform how individuals perceive and relate to sexual experiences, which could lead to an improvement in sexual well-being.

The authors highlight a number of limitations to the study, including the questionnaire-based nature of the data. They also highlight that participants were predominantly white, heterosexual, employed and well-educated, which limits how generalizable the findings are across populations.

The authors caution that while these findings are encouraging, previous trials assessing psilocybin for depression took place under controlled, clinical conditions, using a regulated dose formulated in a laboratory, and involved extensive psychological support before, during and after dosing, provided by mental health professionals. Patients with depression should not attempt to self-medicate with psychedelics, as taking substances in the absence of these careful safeguards may not have a positive outcome.

Barba, added, "It's important to stress our work does not focus on what happens to sexual functioning while people are on psychedelics, and we are not talking about perceived 'sexual performance,' but it does indicate there may be a lasting positive impact on sexual functioning after their [psychedelic](#) experience, which could potentially have impacts on psychological well-being."

Dr. David Erritzoe, clinical director of the Center for Psychedelic Research, said, "This study shines yet more light on the far-reaching effects of psychedelics on an array of psychological functioning. While the findings are indeed interesting, we are still far from a clear clinical application, because psychedelics are yet to be integrated into the medical system. In future, we may be able to see a clinical application, but more research is needed."

More information: Tommaso Barba et al, Psychedelics and sexual functioning: a mixed-methods study, *Scientific Reports* (2024). [DOI: 10.1038/s41598-023-49817-4](#)

Provided by Imperial College London

Citation: Psychedelic experiences linked with improved sexual function (2024, February 7) retrieved 10 May 2024 from <https://medicalxpress.com/news/2024-02-psychedelic-linked-sexual-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--