

## Scientists discover how some advanced breast cancers become resistant to hormone therapy

February 7 2024



Credit: Ahmad Ardity, Pixabay

Researchers have identified why some advanced estrogen receptor (ER) positive breast cancers become resistant to hormone therapy, and have identified drugs currently in development that could keep disease stable for longer for these patients.

A team of scientists at The Institute of Cancer Research, London, found



that for 4% of patients, mutations in the estrogen receptor gene (ESR1) called F404, when combined with specific pre-existing mutations, caused overt resistance to <u>fulvestrant</u> hormone therapy.

They also found that cells with these mutations remained sensitive to a range of compounds which are currently being tested in <u>clinical trials</u>. The researchers hope that in the future, if these new drugs are approved, patients likely to develop <u>treatment resistance</u> through F404 mutations could be identified with a <u>blood test</u> and offered new, <u>alternative</u> <u>treatments</u>.

Fulvestrant is a widely used type of <u>hormone therapy</u> that's usually given to people with ER-positive breast cancer, either as a first line of treatment or once other drugs have stopped working. However, patients' cancers will very often develop resistance to the treatment over time.

The <u>research</u>, which was published in the journal *Cancer Discovery*, involved studying blood samples donated by people taking part in the plasmaMATCH clinical trial.

In the plasmaMATCH clinical trial, managed by the Clinical Trials and Statistics Unit at The Institute of Cancer Research (ICR), researchers are using blood tests—also called liquid biopsies—to analyze small traces of cancer DNA in the blood that have been released from tumor cells.

## Using blood tests to study cancer

The team, working in the Breast Cancer Now Toby Robins Research Center at the ICR, analyzed these blood samples, looking at <u>genetic</u> <u>mutations</u> in the circulating tumor DNA of patients with advanced ERpositive breast cancer and observing how they responded to fulvestrant.

ER-positive breast cancers use estrogen in the body to help them to



grow, and hormone therapies like fulvestrant target the estrogen receptor to prevent this.

Researchers in this study wanted to understand how mutations in the gene that codes for the estrogen receptor, ESR1, can contribute to fulvestrant resistance.

They found that in 4% of patients, following fulvestrant treatment, their breast cancer developed specific mutations in the ESR1 gene, called F404. These new mutations only occurred in patients who already had certain existing mutations in the ESR1 gene before treatment. The researchers noted that the combined effect of these pre-existing and new mutations was a profound resistance to fulvestrant.

## Searching for alternative treatments

The team then tested a series of hormone therapies currently in clinical development on cancer cells with the F404 mutations. They were encouraged to discover that fulvestrant-resistant cancer cells with F404 mutations were sensitive to all four therapies tested.

Professor Nicholas Turner, Professor of Molecular Oncology at The Institute of Cancer Research, London, and Consultant Medical Oncologist at The Royal Marsden NHS Foundation Trust, said, "Our study makes use of innovative blood tests which detect genetic changes present in a patient's cancer, without the need for any invasive procedures.

"The discovery of these specific genetic changes in estrogen receptor positive breast cancer helps to explain one mechanism by which some patients develop resistance to fulvestrant, and which upcoming treatments will be likely to work instead.



"This could revolutionize the way we treat breast cancer, by making use of these simple blood tests to match patients to alternative treatments, bringing them the best possible outcome."

Dr. Kotryna Temicinaite, Head of Research Communication and Engagement at Breast Cancer Now, said, "These findings help us understand how secondary breast cancer can become resistant to hormone therapies like fulvestrant and what other treatments we could use in the future if this resistance happens. With an estimated 61,000 people living with secondary breast cancer in the U.K., research like this is vital."

Dr. Nisharnthi Duggan, Science Engagement Manager at Cancer Research U.K., said, "Research discoveries like this help scientists find better ways to treat cancer. Understanding why a drug like fulvestrant stops being effective will help researchers and doctors identify the best medicines available for people living with certain types of <u>breast cancer</u> and develop new therapies. This would provide people living with specific types of cancer with more effective treatment options, ultimately helping them to live longer, better lives."

**More information:** Belinda Kingston et al, ESR1 F404 Mutations and Acquired Resistance to Fulvestrant in ESR1-Mutant Breast Cancer, *Cancer Discovery* (2023). DOI: 10.1158/2159-8290.CD-22-1387

## Provided by Institute of Cancer Research

Citation: Scientists discover how some advanced breast cancers become resistant to hormone therapy (2024, February 7) retrieved 8 May 2024 from <u>https://medicalxpress.com/news/2024-02-scientists-advanced-breast-cancers-resistant.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.