

Uptick in behavioral disorders reported in children following prenatal exposure to anesthesia

February 29 2024



Credit: Pixabay/CC0 Public Domain

Children prenatally exposed to anesthesia for maternal surgery during pregnancy were significantly more likely than unexposed children to

receive a diagnosis of a disruptive or internalizing behavioral disorder, including attention deficit hyperactivity disorder, behavioral disorders, developmental speech or language disorders, and autism, according to a new study at Columbia University Mailman School of Public Health and Columbia University Irving Medical Center (CUIMC).

Until now, associations between [prenatal exposure](#) to general anesthesia for maternal surgery during pregnancy and subsequent risk of [behavioral disorders](#) in children were unclear. The findings are [published](#) in the *British Journal of Anesthesia*.

Prenatal exposure to general anesthesia was associated with a 31 percent increased risk for a subsequent disruptive or internalizing behavioral disorder (DIBD) diagnosis in children, with higher risk when the exposure occurred in the second or third trimester of pregnancy. The probabilities of DIBD diagnosis between exposed and unexposed children were similar in the first three years of life but diverged thereafter.

"For several years, there have been concerns about the long-term neurodevelopmental effects of anesthetic exposure in children, with the FDA in 2016 releasing a Drug Safety Communication warning against repeated or lengthy use of anesthesia in young children and pregnant women," said Caleb Ing, MD, MS, associate professor of Anesthesiology and Epidemiology at Columbia Public Health and CUIMC and first author.

"Studies evaluating exposures in children, however, have been limited because children needing surgery and anesthesia may have more underlying health problems which may predispose them to [neurodevelopmental disorders](#).

Prenatal exposures are particularly relevant because general anesthetic

drugs cross the placenta, and since the need for anesthesia is related to maternal medical problems, the risk of bias due to underlying health problems in children is reduced, according to the authors. "However, caution is advised, as many procedures in pregnant women may be necessary, and avoidance of necessary procedures can have detrimental effects on mothers and their children," advises Ing.

The study population was drawn from Medicaid Analytic eXPrenatal eXtract for general anesthetic exposure and was identified using maternal claims with International Classification of Disease for either an appendectomy or cholecystectomy which are the two most common non-obstetric procedures performed during pregnancy. Exposures were classified as occurring in the first, second, or third trimester of pregnancy.

The researchers also controlled for a variety of factors including age, race, ethnicity, state of residence, income based on zipcode, and Medicaid enrollment for disability vs. poverty.

Each prenatally exposed child was matched with five unexposed children. The population consisted of 16,778,231 deliveries linked to an infant born between 1999 and 2013 with known sex. Of those, 34,271 children were exposed to general anesthesia due to maternal appendectomy or cholecystectomy during pregnancy.

"This study provides compelling evidence that prenatal exposure to surgery and [general anesthesia](#) may have adverse effects on children's neurobehavioral development ," said Guohua Li, MD, DrPH, professor of Epidemiology and Anesthesiology at Columbia Mailman School and Vagelos College of Physicians and Surgeons, and senior author.

"Our findings could help inform preoperative risk assessment in [pregnant women](#), especially when the surgical procedure is elective or

when viable treatment is available. Results of this study also give added impetus to research on safer anesthetic drugs and techniques."

More information: Caleb Ing et al, Behavioural disorders after prenatal exposure to anaesthesia for maternal surgery, *British Journal of Anaesthesia* (2024). [DOI: 10.1016/j.bja.2024.01.025](https://doi.org/10.1016/j.bja.2024.01.025)

Provided by Columbia University's Mailman School of Public Health

Citation: Uptick in behavioral disorders reported in children following prenatal exposure to anesthesia (2024, February 29) retrieved 28 April 2024 from <https://medicalxpress.com/news/2024-02-uptick-behavioral-disorders-children-prenatal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.