

Vest can detect earlier signs of heart muscle disease

February 26 2024



A volunteer wears the ECGI vest. Middle: Dr George Joy with the vest on a mannequin. Bottom: The team of researchers and staff who have been involved in the vest's use and development. Credit: UCL Institute of Cardiovascular Science / James Tye

A reusable vest that can map the electric impulses of the heart in fine detail could detect abnormalities from a potentially fatal heart disease



much earlier than is currently possible, suggests a new study led by UCL researchers.

The study, <u>published</u> in the *Journal of the American College of Cardiology*, found that an electrocardiographic imaging (ECGI) vest, developed by Dr. Gaby Captur at UCL, could detect electrical changes associated with an inherited heart muscle condition at a stage when standard tests do not pick up signs of disease.

The condition, called <u>hypertrophic cardiomyopathy</u>, is where the muscle wall of the heart becomes thicker and stiffens, affecting how well the heart can pump blood around the body. It affects an estimated one in 300 adults.

While people with genetic variants that cause the disease have no symptoms at all, the condition can lead to heart failure and is frequently cited as the most common cause of sudden, unexpected death in <u>young people</u>.

Lead author Dr. George Joy (UCL Institute of Cardiovascular Science and Barts Heart Centre) said, "By finding subtle electrical abnormalities using our new technique, we are able to detect hypertrophic cardiomyopathy earlier. This is important as it means we can potentially act earlier, providing new treatment to slow the disease as well as fasttracking individuals to clinical trials that have the potential to stop the disease entirely."

"Next steps of the research include repeating these results in a larger group of patients and following individuals over time to see how these early electrical changes affect the risk of life-threatening heart rhythms later on."

The ECGI vest has 256 sensors rather than the 12 used in a standard



electrocardiogram (ECG) and can provide detailed electrical mapping of the heart in just five minutes.

Previously, this kind of detailed mapping was rare—either requiring a catheter to be inserted inside the heart cavity or using single-use devices that were costly and time-consuming to set up. The ECGI vest is also reusable, and so has the potential to be used as a standard screening tool.

The new study looked at 174 patients who had <u>genetic testing</u> (overseen by Dr. Luis Lopes, UCL) recruited from three London hospitals (Barts Heart Centre, St George's Hospital, and Royal Free Hospital), and 37 healthy volunteers. Patients included people who already had hypertrophic cardiomyopathy as well as individuals with disease-causing genetic mutations who did not have overt signs of the disease.

The team found that the ECGI vest identified electrical abnormalities among 1 in 4 individuals with a gene mutation for whom no signs of disease were detected via either <u>cardiac magnetic resonance</u> imaging (MRI), the highest standard of heart imaging, or a 12-lead ECG, the typical way the heart's electrical activity is assessed.

In particular, these patients were found via ECGI to have an uneven pattern of electrical signal recovery and slowed conduction of electrical signals through the heart.

The researchers also applied a machine learning model to the results of 12 markers from the ECGI vest in order to grade the severity of the disease and estimate the risk of sudden cardiac death.

They found this grading matched the risk estimated using a standard protocol, which is based on information such as age and certain structural features of the heart.



Dr. Captur (UCL Institute of Cardiovascular Science and the Royal Free Hospital, London), who was senior author on the latest study, said, "The ECGI vest we have developed is expanding our ability to understand the electrical functioning of the heart and to assess more precisely people's risk of developing life-threatening heart rhythms."

"People who have genetic mutations causing hypertrophic cardiomyopathy are monitored regularly and given advice around exercise. In some cases, this might be to reduce or stop any intense exercise. This prescription can have a huge impact on a person's quality of life, particularly in athletes or young patients. By better understanding risk, we hope to avoid instances where people are given such advice unnecessarily."

Dr. Sonya Babu-Narayan, Associate Medical Director at the British Heart Foundation and consultant cardiologist, said, "The ECG has been at the heart of a doctor's diagnostic toolkit for more than 100 years, allowing a window for us to understand the electrical pulses that control the beat of the heart."

"While further research is needed, this study suggests that the ECGI vest, which collects far more detailed electrical signal data than a routine ECG, could support better identification of high-risk patients with hypertrophic cardiomyopathy. Some people living with hypertrophic cardiomyopathy are particularly susceptible to cardiac arrest. The ECGI vest could offer an opportunity for doctors to better diagnose them early in their disease course and target their treatment to prevent this."

More information: George Joy et al, Electrophysiological Characterization of Subclinical and Overt Hypertrophic Cardiomyopathy by Magnetic Resonance Imaging-Guided Electrocardiography, *Journal of the American College of Cardiology* (2024). DOI: 10.1016/j.jacc.2024.01.006



Provided by University College London

Citation: Vest can detect earlier signs of heart muscle disease (2024, February 26) retrieved 12 May 2024 from <u>https://medicalxpress.com/news/2024-02-vest-earlier-heart-muscle-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.