

Should ADHD medications be stopped during pregnancy?

March 6 2024



Credit: Unsplash/CC0 Public Domain

A new study led by The University of Western Australia looked at whether medications used to treat attention-deficit hyperactivity disorder (ADHD) should be stopped during pregnancy.

Ph.D. student Dani Russell, from UWA's School of Population and Global Health, was the lead author of the paper [published](#) in *Archives of Women's Health*.

"ADHD is becoming more commonly diagnosed in women; consequently, more women are taking ADHD medication such as dexamphetamine during pregnancy," Russell said. "Our study investigated whether there were reasons to stop taking this medication during pregnancy."

The research included women from Western Australia who had been taking dexamphetamine during pregnancy and gave birth between 2003 and 2018.

"The women who stopped taking their medication during pregnancy had an increased risk of threatened pregnancy loss compared to those who continued to take it," Russell said. "However, the results indicated that continuing to take dexamphetamine did not put women or their babies at increased [health risks](#)."

The study found there also appeared to be benefits associated with stopping medication use before conception including decreased risk of pre-eclampsia, hypertension, postpartum hemorrhage, admittance to the special care unit and fetal distress.

"Pregnancy can be a really stressful period for women with ADHD, so it is good to know that stopping their medication during pregnancy may not always be necessary," she said.

The research is part of a larger body of work being carried out with guidance from women in the community with ADHD, hoping to provide much needed insight into the treatment during [pregnancy](#).

More information: Danielle J. Russell et al, Investigating maternal and neonatal health outcomes associated with continuing or ceasing dexamphetamine treatment for women with attention-deficit hyperactivity disorder during pregnancy: a retrospective cohort study, *Archives of Women's Mental Health* (2024). [DOI: 10.1007/s00737-024-01450-4](https://doi.org/10.1007/s00737-024-01450-4)

Provided by University of Western Australia

Citation: Should ADHD medications be stopped during pregnancy? (2024, March 6) retrieved 28 April 2024 from <https://medicalxpress.com/news/2024-03-adhd-medications-pregnancy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.