

## Age and sex associated with patient's likelihood of antimicrobial resistance

March 14 2024



Researchers find that age and sex are associated with patient's likelihood of antimicrobial resistance. Credit: CDC, Unsplash (CC0, creativecommons.org/publicdomain/zero/1.0/)



A person's age, sex and location are correlated with the chance that they have a bloodstream infection that is resistant to antibiotics, according to a new study <u>published</u> in *PLOS Medicine* by Gwenan Knight of the London School of Hygiene and Tropical Medicine, UK, and colleagues.

Antimicrobial resistance (AMR), in which infections cannot be treated with antibiotics, is a major global public health threat. Little has been known about how the prevalence of resistance varies with age and sex even though <u>antibiotic usage</u>, changes in immune function, and exposure to high-risk settings are all linked to age and sex.

In the new study, researchers analyzed data collected as part of routine surveillance between 2015 and 2019 on bloodstream infections in 944,520 individuals across 29 European countries. The team looked at which bacterial species were isolated and sent to the surveillance service, and which antibiotics were used to treat the infections.

Distinct patterns in resistance prevalence by age were observed throughout Europe but varied across bacterial species. For most but not all bacteria, peaks in resistance were seen at the youngest and oldest ages.

The occurrence of methicillin-resistant Staphylococcus aureus (MRSA) increased with age and the occurrence of aminopenicillin resistance in Escherichia coli decreased with age.

Some antimicrobial resistance profiles peaked in middle age; Pseudomonas aeruginosa was most likely to be resistant to several antibiotics around 30 years of age and, for women, the incidence of bloodstream infections due to E. coli peaked between ages 15 and 40. There were other important differences between sexes; in general, men



had a higher risk of antimicrobial resistance than women.

"These findings highlight important gaps in our knowledge of the epidemiology of antimicrobial resistance that are difficult to explain through known patterns of antibiotic exposure and health care contact," the authors say. "Our findings suggest that there may be value in considering interventions to reduce antimicrobial resistance burden that take into account important variations in <u>antimicrobial resistance</u> prevalence with age and sex."

The authors add, "Our findings, that the prevalence of <u>resistance</u> in bloodstream infections across Europe varies substantially by age and sex, highlights important gaps in our knowledge of the spread and selection of AMR."

"In order for us to address this growing threat to public health, we now need data from a wider range of sources to determine the contribution that cultural versus natural history differences have in driving these patterns globally and the role that they play in the increasing rates of AMR being seen."

**More information:** Antimicrobial resistance prevalence in bloodstream infection in 29 European countries by age and sex: An observational study, *PLoS Medicine* (2024). <u>DOI:</u> <u>10.1371/journal.pmed.1004301</u>. journals.plos.org/plosmedicine .... journal.pmed.1004301

## Provided by Public Library of Science

Citation: Age and sex associated with patient's likelihood of antimicrobial resistance (2024, March 14) retrieved 9 May 2024 from <u>https://medicalxpress.com/news/2024-03-age-sex-patient-</u>



likelihood-antimicrobial.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.